

## Integrated use of Cattle manure and Mineral NP on Soil Physico-Chemical Characteristics and Tuber Yield of Potato (*Solanum tuberosum* L.) in Masha District, South-western Ethiopia

Efecto de la aplicación combinada de estiércol de ganado, nitrógeno mineral y fertilizante de fósforo en las características físico-químicas del suelo y el rendimiento de la papa en el tubérculo (*Solanum tuberosum* L.) en el distrito de Masha, suroeste de Etiopía.

Isreal Zewide<sup>1</sup>, Tamado Tana<sup>2</sup>, Lemma Wogi<sup>3</sup> and Ali Mohammed<sup>4</sup>

1- Mizan-Tepi University Department of Horticulture, P.O.Box260, Mizan Teferi, Ethiopia (zewideisreal@gmail.com)

2- Haramaya University, School of Plant Sciences, P.O, Box 138, Dire Dawa, Ethiopia

3- Haramaya University, School of Natural Resources Management and Environmental Sciences, P.O.Box 138, Dire Dawa, Ethiopia

4- Jimma University, Department of Post-harvest Management, P.O.Box 37, Jimma, Ethiopia

### ABSTRACT

Soil fertility depletion due to soil erosion and nutrient leaching caused by heavy and continuous rainfall is one of the major causes for declining of soil productivity and crop yields in Masha district of south-western Ethiopia. Hence, a field experiment was conducted in Belg and Meher seasons at Abelo area, Masha district, south-western Ethiopia, to determine the effect of combined application of cattle manure (CM) with inorganic nitrogen (N) and phosphorus (P) on potato tuber yield, and selected soil physico-chemical characteristics. The treatments comprised of combinations of three rates (2.5, 5, 7.5 t ha<sup>-1</sup>) of CM with 25%, 50% and 75% of recommended rates of mineral NP, respectively. In addition, 100% recommended rate of mineral NP (165 kg N ha<sup>-1</sup> and 137 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup>) and zero rates were used for comparison. The experiment was laid out in a randomized complete block design with three replications. The results revealed that applying 7.5 t ha<sup>-1</sup> CM combined with 75% mineral NP gave significantly highest potato total tuber yield of (40 and 34 t ha<sup>-1</sup>), the same rate increased tuber yield by 55.9% and 43.45% in Belg and by 51.19%, and 36.64% in Meher

over the control and the application of 100% recommended rate of NP fertilizers, respectively soil pH (5.15 and 4.94), Organic carbon (21.1 and 12.7 g kg<sup>-1</sup>), total N (1.71 and 1.51 g kg<sup>-1</sup>), available P (7.55 and 6.27 mg kg<sup>-1</sup>), CEC (22.04 and 23.96 cmol (+) kg<sup>-1</sup>), exchangeable Ca<sup>+2</sup> (8.36 and 8.28 cmol (+) kg<sup>-1</sup>), exchangeable Mg<sup>+2</sup> (3.14 and 2.06 cmol (+) kg<sup>-1</sup>), total porosity (51.1 and 54.97%) more over the treatment decreases exchangeable acidity (2.32 and 2.08 cmol+kg<sup>-1</sup>) and exchangeable aluminum (1.46 and 1.41 cmol(+) kg<sup>-1</sup>), bulk density (1.07 and 1.30 g cm<sup>-3</sup>) in Belg and Meher seasons, respectively. Therefore, it can be concluded that the use of combined application of 7.5 t of ha<sup>-1</sup> CM together with 75% of recommended rates of inorganic N (123.75 kg N ha<sup>-1</sup>) and P (103.05 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup>). can significantly increase potato tuber yield and improve the soil properties and reduces soil acidity.

Keywords: Cattle manure, physico-chemical properties, nitrogen, phosphorus, tuber yield.

#### RESUMEN

El agotamiento de la fertilidad del suelo debido a la erosión del suelo y la lixiviación de nutrientes causada por la lluvia intensa y continua es una de las principales causas de la disminución de la productividad del suelo y los rendimientos de los cultivos en el distrito de Masha, en el suroeste de Etiopía. Por lo tanto, se realizó un experimento de campo en las estaciones de Belg y Meher en el área de Abelo, distrito de Masha, en el sudoeste de Etiopía, para determinar el efecto de la aplicación combinada de estiércol de ganado (CM) con nitrógeno inorgánico (N) y fósforo (P) en Producción de tubérculos de papa y características físico-químicas del suelo seleccionado. Los tratamientos comprendieron combinaciones de tres tasas (2.5, 5, 7.5 t ha<sup>-1</sup>) de CM con 25%, 50% y 75% de las tasas recomendadas de NP mineral, respectivamente. Además, se usaron 100% de la tasa recomendada de mineral NP (165 kg N ha<sup>-1</sup> y 137 kg de P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup>) y tasas cero para la comparación. El experimento se presentó en un diseño de bloques completos al azar con tres repeticiones. Los resultados revelaron que la aplicación de 7.5 t ha<sup>-1</sup> CM combinada con un 75% de NP mineral dio un rendimiento de tubérculo total de papa significativamente más alto (40 y 34 t ha<sup>-1</sup>), la misma tasa aumentó el rendimiento de tubérculo en un 55.9% y 43.45% en Belg y un 51,19%, y un 36,64% en Meher sobre el control y la aplicación del 100% recomendado de fertilizantes NP, respectivamente pH del suelo (5,15 y 4,94), carbono orgánico (21,1 y 12,7 g kg<sup>-1</sup>), N total (1,71 y 1,51 g kg<sup>-1</sup>), P disponible (7.55 y 6.27 mg kg<sup>-1</sup>), CEC (22.04 y 23.96 cmol (+) kg<sup>-1</sup>), Ca + 2 intercambiable (8.36 y 8.28 cmol (+) kg<sup>-1</sup>), intercambiable Mg + 2 (3,14 y 2,06 cmol (+) kg<sup>-1</sup>), porosidad total (51,1 y 54,97%) más sobre el tratamiento disminuye la acidez intercambiable (2,32 y 2,08 cmol + kg<sup>-1</sup>) y aluminio intercambiable (1,46 y 1,41 cmol (+) kg<sup>-1</sup>), densidad aparente (1.07 y 1.30 g cm<sup>-3</sup>) en las estaciones de Belg y Meher,

respectivamente. Por lo tanto, se puede concluir que el uso de la aplicación combinada de 7.5 t de ha<sup>-1</sup> CM junto con el 75% de las tasas recomendadas de N inorgánico (123.75 kg N ha<sup>-1</sup>) y P (103.05 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup>). puede aumentar significativamente el rendimiento del tubérculo de papa y mejorar las propiedades del suelo y reduce la acidez del suelo.

Palabras clave: estiércol de ganado, propiedades físico-químicas, nitrógeno, fósforo, rendimiento de tubérculos.

## INTRODUCTION

Land degradation is a serious and widespread problem for agricultural development and food security in developing countries (Adejumo *et al.*, 2010; Dell'Angelo *et al.*, 2017). The wearing a way of soil in Ethiopian high land is quite pronounced with the average annual soil erosion rate nationwide was estimated at 12 tons ha<sup>-1</sup>, giving total annual soil loss of 1493 million tons (EPA, 2012). As a result, most of the Ethiopian soils, particularly within the highlands, are low in nutrient content due to erosion, leaching and absence of nutrient utilization (Zelege *et al.*, 2010)

The major causes of land degradation in Ethiopia high-land area are natural and anthropogenic factors such as rapid population growth, deforestation, overgrazing, low vegetative cover and unbalanced crop and livestock production. limited recycling of dung and crop residues to the soil, limited application of external sources of plant nutrients, resulting in low nutrient content (Temesgen *et al.*, 2014).

Potato (*Solanum tuberosum* L.) is one of the most important tuber crops in the world. It's tuber consists of mainly carbohydrates, proteins, and lipids. The tuber is used locally alone or together with meat, vegetable and a substituent with a pulse in stew preparation in Sheka Zone, south-western Ethiopia (Isreal *et al.*, 2012). It is a preferable crop in the study area due to that it can be produced more than twice per year. In Masha District, south-western Ethiopia, the potato is one of the major and widely cultivated tuber crops covering an area of 15% of arable land (3255ha) with the total production of 24447 t and productivity of 7.52 t ha<sup>-1</sup> in 2014/15 cropping season (SZARDO, 2015), which is far below the average national yield of 13.5tha<sup>-1</sup> (CSA, 2015).

It is one of the heavy feeders of root and tuber crop requiring relatively large quantities of fertilizers. However, leaching of nutrients due to heavy and continuous rainfall, soil acidity, low nutrient reserves in arable soils and poor crop residue management by potato growers, are the major factors contributing to low soil fertility and low yield of potato in the study area

In addition to the high cost, continuous use of mineral fertilizer is leading to declining soil chemical physical and, biological properties and thus, negatively affecting overall total soil health (Mahajan *et al.*, 2008; Compaore *et al.*, 2011; Bayu and Gebeyehu, 2013)., nutrients supplied exclusively through chemical sources, , lead to unsustainable productivity over the years (Mahajan *et al.*, 2008: Fekadu, 2016). Thus, the undesirable impacts of chemical fertilizers, coupled with their high cost, have prompted the interest in the use of organic fertilizers as a source of nutrients. The combined application of organic together with inorganic fertilizers has been reported to improve crop growth by supplying plant nutrients including micro-nutrients as well as improving soil physical, chemical, and biological properties thereby provide a better environment for root growth by improving the soil structure (Dejene and Lemlem, 2012: Kang *et al.*, 2016).

Major inputs of soil nutrients like the addition of mineral fertilizers, organic sources, and, biological nitrogen fixation are less than major outputs such as harvested products, crop residue removal, leaching, gaseous losses, and soil erosion. This shows a net loss of soil fertility or imbalance between the inputs and the outputs as reported by Lesschen *et al.* (2007) this is also true In Masha district, south-western Ethiopia and encourages the application of organic sources and mineral nitrogen and phosphorus to correct balances of nutrient inputs and outputs.

The combined application of organic and inorganic fertilizers can improve soil health and nutrients supply for plant growth and subsequently would be resulted in economic profitability. Many research findings showed that neither inorganic fertilizers nor organic sources alone can result in the sustainable productivity (Yadav *et al.*, 2013: Fekadu, 2016). Furthermore, the price of inorganic fertilizers is increasing and becoming unaffordable for resource-poor smallholder farmers. The best remedy for soil fertility management is, therefore, a combination of both inorganic and organic fertilizers, where the inorganic fertilizer provides readily available nutrients and the organic fertilizer mainly increases soil organic matter and improves soil structure and buffering capacity (Yadav *et al.*, 2017). The combined application of inorganic and organic fertilizers, usually termed as integrated nutrient management, is widely recognized as a way of increasing crop yields through sustainable soil health and productivity (Mahajan *et al.*, 2008; Kang *et al.*, 2016). Several researchers (*e.g.* Mahajan *et al.*, 2008; Getachew, 2009; Gafar *et al.*, 2012: Agegnehu and Amede, 2017) verified the beneficial effect of integrated nutrient management in moderating the deficiency of a number of macro- and micro-nutrients.

In view of this fact, identifying the optimum dose of integrated organic and inorganic fertilizers is crucial and is required to supply sufficient amount of nutrients for increased crop yields. Daniel and Nigussie (2006) reported high tuber yield of potato (29.59 t ha<sup>-1</sup>) when cattle manure at the rate of 10tha<sup>-1</sup> was combined with inorganic nitrogen at 111 kg Nha<sup>-1</sup> and

phosphorus at 90 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> on Nitosols. Similarly, Shiferaw (2014) reported the highest mean potato tuber yield (41.88 t ha<sup>-1</sup>) with combined application of 15t ha<sup>-1</sup>cattle manure with the application of 100% recommended rate of NP (100kg ha<sup>-1</sup>), which increased tuber yield by 393.9%, over control. Likewise, Biruk *et al.* (2015) obtained the highest potato tuber yield (38.68 t ha<sup>-1</sup>) by application of 30 t ha<sup>-1</sup>of cattle manure along with nitrogen at 120kg N ha<sup>-1</sup> and phosphorus at 92 kg P<sub>2</sub>O<sub>5</sub>ha<sup>-1</sup>. Thus, the integrated nutrient management involving judicious use of combined organic and inorganic fertilizer sources is a feasible and productive approach to overcome soil fertility constraints (Efthimiadou *et al.*, 2010).

However, research on integrated nutrient management for potato production has not yet been conducted at Abelo area, Masha district of Sheka Zone, south-western Ethiopia where soil fertility depletion is very severed. In order to address the key issue of soil fertility depletion attributed and reduction of potato yields the study area, different rates of cattle manure and mineral NP was conducted in Belg and Meher seasons. The objective of the study was to determine effects of combined application of CM with inorganic NP fertilizers on selected soil physico-chemical properties and potato tuber yield and subsequently to improve soil fertility and crop productivity of the study area.

## MATERIALS AND METHODS

Description of the Study Site: The experimental site lies between 7°44'17.6"- 7°44'20.1" N latitude, 35°29'18"- 35°29'41.9" E longitude and located at an elevation of 2205 m.a.s.l. The total area of the sites 5603m<sup>2</sup> (0.56 ha), which is found in Abelo area, Masha District of Sheka Zone in South-west Ethiopia (Figure 1). Abelo area is a well-known area of high potential for root and tuber crops production including potato in Masha District.

The rainfall pattern of the site is characterized by monomodal distribution with the relatively small rainy season in Belg (February -May) and main rainy season's Meher (June - October). The total rainfall was 633 and 1153 mm in 2016 Belg and Meher cropping seasons, respectively.

The maximum temperature of 2016 *Belg* season varied from 24.98 °C (February) to 24.62 (May) and 24.74 (June) to 26.19 °C (October) and mean maximum temperature of 25.05 and 25.56 °C in *Belg* and *Meher* seasons, respectively. The minimum temperature varied between 16.01°C (February) to 13.9°C (May) in *Belg* and 12.64(June) to 13.48°C (October) in the *Meher* season. Mean minimum temperature was 14.5 and 13.15°C in *Belg* and *Meher* seasons, respectively as indicated in Figure 2.




Figure 1. Map of Ethiopia showing the location of SNNPRS (A), a map of SNNPRS showing the location of the study area (B) and a map of the study area showing the location of soil sampling points




Figure 2. Monthly total rainfall (mm) and mean monthly minimum (Tmin) and maximum (Tmax) temperatures (°C) of the study site in 2016 Belg(A) and 2016 Meher(B) cropping seasons. Source: Masha Meteorological Station (2016)

Experimental Materials: potato variety 'Belete' from Holeta Agricultural Research Center was used for this experiment. The variety was released in 2009 and has been accepted by farmers due to its high yielding ability, consumers' preference, wider adaptation, better cooking ability and relative resistance to late blight compared to local and other improved potato varieties. As a result, this variety is being cultivated widely in the study area. It is well performing at an altitude of 1600-2800m.a.s.l and requires rainfall of 750-1000 mm per season. It reaches maturity within 90-120 days. The productivity of this variety ranges from 29.13 t ha<sup>-1</sup> under farmers' field and 44.8 t ha<sup>-1</sup> at research stations (MOARD, 2012).

Urea (46% N) and TSP (46% P<sub>2</sub>O<sub>5</sub>) were used as inorganic N and P sources whereas CM was used as an organic fertilizer. CM was obtained from sustainable land management (SLM) project in Masha district; where Urea and TSP were obtained from Teppi Soil Testing Research Center.

The pH and electrical conductivity (EC) of the cattle manure were determined in extracts of 1:10(w/v) cattle manure to distilled water ratio as described by Ndegwa and Thompson (2001). Total N was measured by the Kjeldahl method. Organic carbon (C) was determined by the Walkley and Black method (Walkley and Black 1934). Total P, Ca, Mg and K were extracted following the wet digestion method (Okalebo *et al.*, 2002). The concentration of P was determined by the molybdenum blue coloration method. Total cations (Ca and Mg) were determined from the wet digested samples by atomic absorption spectrophotometer, while total K and Na were estimated by a flame photometer (Okalebo *et al.*, 2002). The results of all analyzed parameters for cattle manure are depicted in Table 3.

Treatments and Experimental Design: treatments consisted of three rates of cattle manure (2.5, 5, 7.5 t ha<sup>-1</sup>) and three rates of recommended inorganic NP fertilizers (0, 25%, 50%, and 75%) and 100% recommended rate of inorganic NP fertilizer (165 kg N ha<sup>-1</sup> and 60 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup>) and zero rates were used. The field experiment was laid out in the Randomized Complete Block Design (RCBD) with three replications. Gross and net plot sizes were (4.5m row width \* 3.6m row length) 16.2 m<sup>2</sup> and (3m row length\*3m row width) 9m<sup>2</sup>, respectively. Treatments were assigned to each plot randomly. Each plot was comprised of six rows with 12 plants/hills. The total number of plants per plot was 72. Spacing was 30 cm between plants/hills within rows and 75 cm between rows, which gave a planting density of 4.4 plants/hills m<sup>-2</sup>. The spacing between plots and replications were 1 and 1.5 m, respectively.

Experimental Procedure: the experimental land was well prepared prior to planting. In the first season, it was plowed three times with oxen and the plots were leveled manually. The soil was pulverized, leveled using human labor, each individually. The plot was prepared and addition of ridging and then a division of the experimental units was done manually. Cattle

manure was applied to the respective plots thoroughly and then incorporated into the soil layer (0-20 cm) a month prior to planting at the rate of 2.5, 5 and 7.5 t ha<sup>-1</sup> on dry weight bases. Phosphorus was applied in the form of triple superphosphate (46% P<sub>2</sub>O<sub>5</sub>) by drilling along the furrows and per hill and mixed well with the soil. Pre-sprouted potato tuber seeds were planted on 2<sup>nd</sup> February 2016 during *Belg* and on 2<sup>nd</sup> June 2016 during *Meher* seasons. Nitrogen was side-dressed in three split application methods (i.e., 1/4<sup>th</sup> at plant emergence; 2/4<sup>th</sup> at mid-stage-at 40 days after planting; and 1/4<sup>th</sup> at the initiation of tubers) because of its mobility and possible injury to the crop. Conventional agronomic practices such as hoeing, weeding, and earthing-up were carried out uniformly to all plots. Dehaluming were done to thicken tuber periderm when the plants reached physiological maturity and senesced. This was done two weeks before harvesting to toughen the periderm in order to reduce bruising and skinning during harvesting and post-harvest handling. Tubers were harvested on May 15, 2016, in *Belg* and on October 8, 2016, in *Meher* seasons, when 70% of haulms were dried from plants grown in the four middle rows. Harvesting was done manually using hand hoe with great care and only parts of plants grown in the central parts of the plots were used for data collection to reduce border effect. Thus, 40 plants in the net plot were used to collect potato yield and finally converted into tha<sup>-1</sup>.

**Soil Sampling and Analysis:** Prior to planting, soil samples (0 - 20 cm), from ten spots across the experimental fields were collected in a zig-zag pattern, in 2016 during *Belg* and *Meher* cropping seasons. One composite sample was made. The soil sample was prepared for physicochemical analyses. Soil and were air dried ground and passed through 2 mm sieve for the analysis of physico-chemical properties except for total nitrogen and organic carbon which were passed through 0.5 mm sieve.

Soil Particle size distribution was analyzed by the hydrometer method (Day, 1965) after destroying OM using hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>) and dispersing the soils with sodium hexametaphosphate Na(PO<sub>3</sub>)<sub>6</sub>. Soil bulk density was determined by the undisturbed core sampling method (Blake, 1965) after drying the soil samples in an oven at 105 °C to constant weights, while particle density was estimated by the pycnometer method (Barauah and Barthakulh, 1997). Percent total pore space was computed from the values of bulk density (BD) and particle density (PD) as described by Hillel (2004).

$$TP(\%) = [1 - (\frac{BD}{PD})] * 100$$

Soil pH-H<sub>2</sub>O was measured using a pH meter in a suspension of 1:2.5 soils: water ratio. The soil OC content was determined following the wet oxidation method as outlined by Walkley and Black (1934) Total N content in the soil samples was determined titrimetrically following the Kjeldahl method as described by Jackson (1958). Available P was determined using Bray II



method (Bray and Kurtz, 1945). Cation exchange capacity (CEC) and exchangeable bases (Ca, Mg, K and Na) were determined after extracting the soil samples by ammonium acetate (1N NH<sub>4</sub>OAc) at pH 7.0. Exchangeable Ca and Mg in the extracts was analyzed using atomic absorption spectrophotometer while Na and K were analyzed by a flame photometer (Rowell, 1994). Cation exchange capacity was thereafter estimated titrimetrically by distillation of ammonium that was displaced by sodium after leaching with NaCl solution (Chapman, 1965). Percentage base saturation (PBS) was calculated as the ratio of the sum of the charge equivalents of the base-forming cations (Ca, Mg, Na and K) to the CEC of the soil.

Exchangeable acidity was determined by saturating the soil samples with 1M KCl solution and titrated with 0.02 M NaOH as described by Rowell (1994). From the same extract, exchangeable Al in the soil samples was titrated with a standard solution of 0.02 M HCl. Finally, exchangeable H was obtained by subtracting exchangeable Al from exchangeable acidity (Al + H) (Rowell, 1994). Available micronutrients (Fe, Mn, Zn and Cu) were extracted from the soil samples with Diethylene Triamine Pentaacetic Acid (DTPA) as described by Lindsay and Norvell (1978) and all were measured by atomic absorption spectrophotometer at their respective wavelengths.

Just after harvesting of the crop, composite surface (0 - 20 cm) soil samples were collected from three spots for each plot from every replication. These samples were composited to yield one representative sample per replication from each plot for determination of CEC, pH, total N, available P, available K and organic carbon contents using procedures indicated for pre-sowing soil analysis. The extract of K was analyzed using flame photometer (Black, 1965). The bulk density (BD) of the soil was measured from the undisturbed soil samples collected from each plot using core sampler, which was weighed at field moisture, and after drying the pre-weighed core soil sample to a steady weight in an oven at 105 °C according to the procedure described by Okalebo *et al.* (2002), while particle density (PD) was measured using pycnometer (Barauah and Barthakulh, 1997).

$$TP(\%) = [1 - (\frac{BD}{PD})] * 100$$

where, BD = bulk density; PD= particle density (Hillel, 2004).

Statistical Data Analysis: The post-harvest soil and tuber yield data were subjected to analysis of variance (GLM procedure) using SAS software program version 9.2 (SAS Institute, 2003). Homogeneity of variances was calculated using the F-test as described by Gomez and Gomez (1984) and since the F-test has shown heterogeneity of the variances of the two seasons for all of the agronomic parameters, a separate analysis was used for the two

seasons. The Fisher's protected least significant difference (LSD) test at 0.05 probability level was employed to separate treatment means where significant treatment differences occurred.

## RESULTS AND DISCUSSION

Soil physical and chemical properties before planting: The bulk density value of the soil was  $1.37 \text{ g cm}^{-3}$  in Belg and  $1.38 \text{ g cm}^{-3}$  in Meher season where particle density was  $2.58 \text{ g cm}^{-3}$  of soil in Belg and  $2.6 \text{ g cm}^{-3}$  in Meher and total porosity was 46.80% in Belg and 46.92% in Meher. According to Baruah and Barthakulh (1997), soil bulk density and particle density rating of this soil has moderate compaction. The moderate compaction of the soil could be due to continuous cultivation which causes disintegration of the soil structure and exposure of surface soil to the direct raindrops. The total porosity of this soil was 46.96% which was classified as high according to the rating of Hillel (2004). Therefore, the soil is good for optimum movement of air and water through the soil and there is no problem on tuber growth and expansion in the soil. Soil particle size analysis results showed that texture of the soil is sandy clay loam in both seasons with particle size distribution of 57% sand, 18% silt, 25% clay in Belg and 56% sand, 16% silt and 28% clay in Meher seasons (Table 1). This soil is, therefore, expected to have poor structure, unstable because of low content of clay and organic matter, slightly weak in strength when relatively dry, which creates no problem for emerging seedlings and susceptible to leaching leading to loss of available plant nutrients especially basic cations that cause deficient in macronutrients like N, P, K, Ca, Mg and S. These results of the initial soil test analysis showed that the soil at the site was strongly acidic, with pH of (5.01) in Belg and Very Strongly acidic (4.8) in Meher season. Low status of total N of 1 and  $0.8 \text{ (g kg}^{-1}\text{)}$  content could possibly be due to either low initial organic matter content of the soil, which contributes about 90-95% of soil nitrogen or leaching of nitrate by torrential rainfall prevalent in the experimental areas (Aziz *et al.*, 2010; Ahmad and Arshad, 2015). Low status of organic carbon, ( $12 \text{ g kg}^{-1}$ ) in Belg and ( $10.2 \text{ g kg}^{-1}$ ) in Meher seasons, Low status of available phosphorus of the soils 5.5 ppm in Belg and 5 ppm in Meher season. This could be attributed to the poor management of crop residue, poor land management practices, slash-and-burn agricultural systems, low soil organic matter, high soil erosion. low inherent fertility of soil, fixation of phosphorus in acidic soil resulting in nutrient reduction and the decline in soil fertility. Thus, the response to added organic and mineral fertilizer at different season is expected to show responses on the crop and the soil. The soils analysis also revealed that in both study seasons have medium CEC of  $20 \text{ cmol (+) kg}^{-1}$  in Belg season and  $19.3 \text{ cmol (+) kg}^{-1}$ , in Meher season, high in micronutrient cations Fe, Mn, Cu, and Zn both in Belg and Meher season. due to increased availability at low pH, cations being less strongly bound to the soil and readily exchangeable and trace in exchangeable sodium according to the rating of Tekalign *et al.* (1991) (Table 2)

The soil had also relatively high content of exchangeable acidity ( $3.83\text{cmolc}(+) \text{kg}^{-1}$ ) and aluminum ( $3.82\text{cmolc}(+) \text{kg}^{-1}$ ) in Belg and exchangeable aluminum ( $2.01 \text{cmolc}(+)\text{kg}^{-1}$ ) in Belg and ( $2.46 \text{cmolc}(+) \text{kg}^{-1}$ ). in Meher season.

Analysis of the composition of soil and cattle manure revealed better nutrient composition in Belg than in Meher season because in Belg season there is optimum soil temperature, moisture and pH that might create suitable condition for microorganism multiplication both in number and species causes fast decomposition of materials this results in narrow carbon to nitrogen ratio and release of more nutrients Therefore, the microbial-based indicators can be used as early indicators of soil degradation or improvement (Table 1 and 2).

Table 1.Initial soil physical characteristics of the experimental site. During Belg and Meher seasons

Soil parameters	Belg season		Meher season		References
	Value	Rating	Value	Rating	
% Sand	57		56		
% Silt	18		16		
%clay	25		28		
Textural class	Sandy clay loam		Sandy clay loam		
Bd ( $\text{g cm}^{-3}$ )	1.37	Medium	1.38	Medium	Barauahand Barthakulh(1997)
PD ( $\text{g cm}^{-3}$ )	2.58	Medium	2.6	Medium	Barauah and Barthakulh (1997)
%porosity	46.80		46.92	high	Hillel (2004)

Bd=bulk density Pd-particle density

Table 2. Initial soil chemical characteristics of the experimental site. During Belg and Meher seasons

Soil parameters	Belg season		Meher season		
	Value	Rating	Value	Rating	
pH (1:2.5 H <sub>2</sub> O)	5.01	Strongly acidic	4.8	Very strongly acidic	Tekalign <i>et al.</i> (1991)
O.C(g kg <sup>-1</sup> )	12	Low	1.02	Low	Tekalign <i>et al.</i> (1991)
N (g kg <sup>-1</sup> )	1	Low	0.08	Low	Tekalign <i>et al.</i> (1991)
C: N	12	Low	12.75	low	Hazelton and Murphy (2007)
Available P (mg kg <sup>-1</sup> )	5.5	Low	5	Low	Jones and Benton (2003)
Exchangeable Ca (cmol (+) kg <sup>-1</sup> soil)	6.5	Medium	6.3	Medium	FAO (2006)
Exchangeable Mg (cmol (+) kg <sup>-1</sup> soil)	2.1	Moderate	1.4	Moderate	FAO (2006)
Exchangeable K (cmol (+) kg <sup>-1</sup> soil)	0.42	High	0.36	High	FAO (2006)
Exchangeable Na (cmol (+) kg <sup>-1</sup> soil)	0.06	Very low	Nil	Very low	Landon (1991)
CEC (cmol (+) kg <sup>-1</sup> soil)	20	Medium	19.3	Medium	Hazelton and Murphy (2007)
Pbs (%)	45.4	Medium	41.7	Medium	Hazelton and Murphy (2007)
Available potassium (ppm)	96			85	
Exchangeable Al (cmol (+) kg <sup>-1</sup> soil)	2.01	High	2.46	High	Hazelton and Murphy (2007)
Exchangeable acidity (cmol (+) kg <sup>-1</sup> )	3.83	High	3.82	High	Hazelton and Murphy (2007))
Cu (mg kg <sup>-1</sup> ) (DTPA)	8	High	6	High	Jones and Benton (2003)
Fe (mg kg <sup>-1</sup> ) (DTPA)	120	High	80	High	Jones and Benton (2003)
Zn (mg kg <sup>-1</sup> ) (DTPA)	1.5	High	1.2	High	Jones and Benton (2003)
Mn(mgkg <sup>-1</sup> ) (DTPA)	25	High	20	High	Jones and Benton (2003)

DTPA = Diethylene Triamine Pentaacetic Acid, Pbs=percent base saturation

The analysis of cattle manure presented in Table 3 indicated that it contains relatively higher content of carbon, N, P, pH, total N, Ca, Mg, K, Na, CEC, EC and moisture content in Belg season as compared to Meher season possibly Because of its fast decomposition elevated

contents of salts it can be utilized to raise the pH of acid soils Therefore, cattle manure can be used as an alternative to lime either by itself or as a mixture of mineral NP

Table 3. nutrient content of cattle manure used in the experiment in Belg and Meher season

Parameters	Belg	Meher
pH (1:2.5 H <sub>2</sub> O)	6.8	7.1
Electrical conductivity (uS cm <sup>-1</sup> )	182	148
Carbon (g kg <sup>-1</sup> )	250	220
Total N (g kg <sup>-1</sup> )	192	160
C: N	13:1	14;1
Totoal phosphorus (g kg <sup>-1</sup> ))	13.74	119
Total Ca (g kg <sup>-1</sup> )	78.54	67.48
Total Mg (g kg <sup>-1</sup> )	13.3	9.9
Total K (g kg <sup>-1</sup> )	46.74	52.2
Total Na (g kg <sup>-1</sup> )	0.7	0.9
Moisture content (%)	78	1

Mean square value due to combined application of cattle manure and mineral Were highly significant ( $p < 0.01$ ) and low coefficient of variation indicating good precision of the experiment for all the 12 studied soil physicochemical characteristics and tuber yield of potato at Masha in 2016 Belg and Meher season the overall result showed the existence of sufficient variation for studied tuber yield and soil physicochemical properties and result is presented in Table 4.

Table 4. Analysis of variance for potato tuber yield and soil physico-chemical properties after crop harvest.

Soil parameter and tuber yield	Season	Source of variation with a degree of freedom			
		Replication (2)	Treatment (10)	Error (20)	CV (%)
potato tuber yield	Belg	16.43 <sup>ns</sup>	70.09 <sup>**</sup>	4.92	6.74
	Meher	5.75 <sup>ns</sup>	33.28 <sup>**</sup>	1.77	4.62
Bulk density (g cm <sup>-3</sup> )	Belg	0.02 <sup>ns</sup>	0.011 <sup>**</sup>	0.00096	2.7
	Meher	0.017 <sup>ns</sup>	0.00587 <sup>**</sup>	0.0008	2.02
Particle density (g cm <sup>-3</sup> )	Belg	0.00117 <sup>ns</sup>	0.0034 <sup>ns</sup>	0.00095	1.16
	Meher	0.032 <sup>ns</sup>	0.11 <sup>**</sup>	3.22	1.04
Porosity (%)	Belg	0.68 <sup>ns</sup>	2.73 <sup>**</sup>	0.581	1.54
	Meher	36.70 <sup>ns</sup>	5.23 <sup>**</sup>	0.45	1.37
pH (1:2.5 H <sub>2</sub> O)	Belg	0.10 <sup>ns</sup>	0.01 <sup>**</sup>	0.00075	0.54
	Meher	0.02 <sup>ns</sup>	0.007 <sup>**</sup>	0.00049	0.45
O.C	Belg	0.0104 <sup>ns</sup>	0.018 <sup>**</sup>	0.0071508	4.334051
	Meher	0.0026 <sup>ns</sup>	0.017 <sup>**</sup>	0.00333038	5.21
T. N	Belg	0.00014 <sup>ns</sup>	0.00054 <sup>**</sup>	0.00015727	8.672782
	Meher	0.00459 <sup>ns</sup>	0.0012 <sup>**</sup>	0.00003363	4.92
Available phosphorus	Belg	0.61 <sup>ns</sup>	1.19 <sup>**</sup>	0.31837543	8.610600
	Meher	0.099 <sup>ns</sup>	0.315 <sup>**</sup>	0.04958257	3.92
Calcium(cmol(+) kg <sup>-1</sup> )	Belg	0.089 <sup>ns</sup>	1.75 <sup>**</sup>	0.11	4.86
	Meher	0.90 <sup>ns</sup>	0.157 <sup>**</sup>	0.098	2.15
Magnesium(cmol(+) kg <sup>-1</sup> )	Belg	0.375 <sup>ns</sup>	0.16 <sup>**</sup>	0.038	6.79
	Meher	0.23 <sup>ns</sup>	0.035 <sup>**</sup>	0.018	4.62
Potassium(cmol(+) kg <sup>-1</sup> )	Belg	0.00022 <sup>ns</sup>	0.0033 <sup>**</sup>	0.0005	6.17
	Meher	0.000055 <sup>ns</sup>	0.0025 <sup>**</sup>	0.00017	6.18
Sodium (cmol(+) kg <sup>-1</sup> )	Belg	0.000075 <sup>ns</sup>	0.0000051	0.001	23.12
	Meher	0.0029 <sup>ns</sup>	0.0003	35.79	35.79
Exchangeable acidity(cmol(+) kg <sup>-1</sup> )	Belg	0.0096 <sup>ns</sup>	0.63 <sup>**</sup>	0.01	3.63
	Meher	0.02 <sup>ns</sup>	0.09 <sup>**</sup>	0.02	4.86
Exchangeable Aluminum (cmol(+) kg <sup>-1</sup> )	Belg	0.0044 <sup>ns</sup>	0.28 <sup>**</sup>	0.013	5.93
	Meher	0.18 <sup>ns</sup>	0.033 <sup>**</sup>	0.01	4.00
Cation exchange capacity(cmol(+) kg <sup>-1</sup> )	Belg	0.075 <sup>ns</sup>	9.83 <sup>**</sup>	0.63	4.29
	Meher	27.06 <sup>ns</sup>	2.96 <sup>**</sup>	4.10	0.63
Percent base saturation (PBS) (%)	Belg	2.013 <sup>ns</sup>	3.4 <sup>**</sup>	0.19	0.9
	Meher	0.00185 <sup>ns</sup>	0.00185 <sup>**</sup>	0.002	0.38

Ns non-significant more than 5% level of significance ( $P > 0.05$ ), Significant at 5% level of significance ( $P < 0.05$ ) and \*\* highly Significant at 1% level of significance. ( $P < 0.01$ ) as established by LSD test; \*

**Total Tuber Yield of Potato:** Total tuber yield of potato was highly significantly ( $P < 0.01$ ) affected by the application of cattle manure and inorganic NP (Table 4). Increasing the application rates of cattle manure and inorganic NP from 0 to 7.5 t ha<sup>-1</sup> increased the total tuber yield from 25.78 to 40.20 t ha<sup>-1</sup> and 22.63- 34.22 t ha<sup>-1</sup> (Table 5). The highest yield was obtained at 7.5 t ha<sup>-1</sup> CM+75% NP ha<sup>-1</sup>, but the lowest yield was obtained at zero dosage of cattle manure and inorganic NP application. Increasing the application rates of cattle manure and inorganic NP from zero to 7.5 t CM+75% NP ha<sup>-1</sup> increased total tuber yield by 55.9%, 42.90% and 51.19%, 36.64% in both Belg and Meher season as compared to zero and 100% inorganic NP application of fertilizers, respectively. This show there is an opportunity for additional gain in tuber yield through the further application of more rates of combined cattle manure with inorganic NP fertilizers above 7.5 t ha<sup>-1</sup> CM+75% of recommended NP ha<sup>-1</sup>. This result is partly in line with the finding of and Najm *et al.* (2012) reported that combined application of cattle manure and inorganic fertilizers increased the total tuber yield. In his study highest tuber yield (36.8 tha<sup>-1</sup>) was obtained by application of 20 tons<sup>-1</sup> Cattle manure+150 kg N ha<sup>-1</sup>. on clay loam soils of Islamic Azad University, Roudehen Branch, Tehran, Iran, similarly Alemi (2012) reported the maximum tuber yield (53.4 and 67 t ha<sup>-1</sup>) and (51 and 63.1 t ha<sup>-1</sup>) in 2003 and 2004 cropping season in *Nitosol* and *Vertisol*, soils of in addition maximum tuber dry weight (632.38 g·m<sup>-2</sup>) was obtained by integrated use of One third of Crop residue + one third of CM + Bio-fertilizer +150% RDF. In 2012 2013 cropping season the gangetic alluvium belt of West Bengal Shubhadip *et al.* (2017) and also Kabira (2017) integrated the use of cattle manure at rate 2.47 t ha<sup>-1</sup> with mineral NPK (23:23:0) at 77kg ha<sup>-1</sup> and Desiree and Kenya Mavuno potato varieties. in the second season April 2014 to August 2014 at kalroTigoni.station Nairobi Kenya provide better yield (26.24 t ha<sup>-1</sup>).

Results have shown that all the treatments increased total tuber yield in both Belg and Meher season as compared to control, but the increment in total tuber yield is higher in belg season. This may be due to rainfall and temperature variability in each season causes soil fertility dynamics by facilitating more soil organic matter and manure decomposition this may favor higher nutrient availability in Belg season as compared to Meher season. in addition, compatibility of essential nutrient availability with growing phases that is maximum demand at tuber bulking phase and low in early and maturity phase.

Table 5. Tuber yield ( $t\ ha^{-1}$ ) and percent yield change (%) as influenced by the combined application of organic fertilizer and mineral NP in Belg and Meher seasons, in Masha district, south-western Ethiopia

Treatment*	Belg		Meher	
	Total tuber yield ( $t\ ha^{-1}$ )	% change over 100% NP	Total tuber yield ( $t\ ha^{-1}$ )	% change over 100% NP
2.5 $t\ ha^{-1}$ CM+25% NP	29.3 <sup>de</sup>	4.13	26.54 <sup>de</sup>	5.31
2.5 $t\ ha^{-1}$ CM+50% NP	29.9 <sup>d</sup>	6.26	27.98 <sup>d</sup>	10.43
2.5 $t\ ha^{-1}$ CM+75% NP	30.29 <sup>d</sup>	7.65	28.25 <sup>d</sup>	11.39
5 $t\ ha^{-1}$ CM+25% NP	30.98 <sup>d</sup>	10.10	28.31 <sup>d</sup>	11.60
5 $t\ ha^{-1}$ CM+50% NP	35.06 <sup>c</sup>	24.60	28.73 <sup>cd</sup>	13.10
5 $t\ ha^{-1}$ CM+75% NP	35.99 <sup>bc</sup>	27.90	30.88 <sup>bc</sup>	20.74
7.5 $t\ ha^{-1}$ CM+25% NP	36.77 <sup>abc</sup>	30.68	31.13 <sup>b</sup>	21.63
7.5 $t\ ha^{-1}$ CM+50% NP	39.6 <sup>ab</sup>	40.73	32.46 <sup>ab</sup>	26.35
7.5 $t\ ha^{-1}$ CM+75% NP	40.2 <sup>a</sup>	42.87	34.22 <sup>a</sup>	32.61
100% NP	28.14 <sup>de</sup>	0	25.04 <sup>e</sup>	0
Control	25.78 <sup>e</sup>	-8.37	22.63 <sup>f</sup>	-8.57
LSD (5 %)	3.78		2.26	
Significance	**		**	
CV (%)	8.77		4.62	

Means followed by the same letter within a column are non-significantly different to each other at \* Significant at 5% level of significance ( $P < 0.05$ ) and \*\* highly Significant at 1% level of significance. ( $P < 0.01$ ) as established by LSD test; \* CM = Cattle manure in  $t\ ha^{-1}$

Soil Physical Characteristics: The combined application of cattle manure and inorganic N and P fertilizers had highly significant ( $P < 0.01$ ) effect on the bulk density and porosity of the soil both in Belg and Meher season. (Table 6). The lowest bulk density ( $1.28\ g\ cm^{-3}$ ) was obtained from the application rate of  $7.5\ t\ ha^{-1}$  CM in combination with 75% inorganic N and P in Belg season which is stastically the same as application  $7.5\ t\ ha^{-1}$  CM with 50% inorganic N and P,  $7.5\ t\ ha^{-1}$  CM with 25% inorganic N and P and  $5\ t\ ha^{-1}$  CM in combination with 75% inorganic N and P and  $5\ t\ ha^{-1}$  CM in combination with 50% inorganic N and P in Belg season. Similarly, the lowest bulk density ( $1.30\ g\ cm^{-3}$ ) in Meher was obtained from the application rate of  $7.5\ t\ ha^{-1}$  CM in combination with 75% inorganic N and P which is statistically at par with application of  $7.5\ t\ ha^{-1}$  CM in combination with 75% inorganic N and P while highest bulk densities ( $1.25\ g\ cm^{-3}$ ) in Belg and ( $1.38\ g\ cm^{-3}$ ) in Meher seasons were from the control, application of 100% NP and  $2.5\ t\ ha^{-1}$  CM with 25% NP in both Belg and Meher season respectively. (Table 4). The lowest bulk density of  $7.5\ t\ ha^{-1}$  CM applied plots was due to the availability of organic matter to the soil at the proper time and in the proper proportions and this improves soil physical and chemical properties due to increased soil organic matter. These results corroborate with the findings of Paul et al. (2016) reported that cattle manure at  $10\ t\ ha^{-1}$  with 100% RDF ( $N_{150}P_{75}K_{75}$ ) has advantages in improving the soil organic matter and water holding capacity, with an associated decrease in bulk density and increasing infiltration rate of water. Similarly, Fekadu (2016); and Saha et al. (2008) reported that the use of cattle



manure with mineral fertilizer significantly increased soil organic carbon (SOC) and decreased bulk density over time.

Table 6. Bulk density ( $\text{g cm}^{-3}$ ), particle density ( $\text{g cm}^{-3}$ ), and total porosity (%) of soil after crop harvest as influenced by the combined application of cattle manure and mineral NP in Belg and Meher seasons, in Masha district, South-western Ethiopia.

	Belg			Meher		
	Bulk Density	Particle Density	Total Porosity	Bulk Density	Particle density	Total Porosity
T3	1.37 <sup>ab</sup>	2.66	48.55 <sup>ef</sup>	1.37 <sup>ab</sup>	2.67	48.57 <sup>e</sup>
T4	1.36 <sup>bc</sup>	2.65	48.62 <sup>ef</sup>	1.36 <sup>bc</sup>	2.67	48.49 <sup>e</sup>
T5	1.35 <sup>cd</sup>	2.65	49.11 <sup>de</sup>	1.35 <sup>cd</sup>	2.67	48.86 <sup>de</sup>
T6	1.35 <sup>cd</sup>	2.64	49.12 <sup>de</sup>	1.34 <sup>d</sup>	2.67	48.94 <sup>de</sup>
T7	1.33 <sup>e</sup>	2.65	49.63 <sup>d</sup>	1.33 <sup>e</sup>	2.67	49.38 <sup>d</sup>
T8	1.31 <sup>f</sup>	2.65	50.39 <sup>c</sup>	1.31 <sup>f</sup>	2.67	50.06 <sup>c</sup>
T9	1.3 <sup>fg</sup>	2.63	50.58 <sup>bc</sup>	1.30 <sup>fg</sup>	2.62	50.33 <sup>bc</sup>
T10	1.29 <sup>gh</sup>	2.64	51.11 <sup>ab</sup>	1.29 <sup>gh</sup>	2.63	50.85 <sup>ab</sup>
T11	1.28 <sup>h</sup>	2.65	51.79 <sup>a</sup>	1.30 <sup>h</sup>	2.61	51.10 <sup>a</sup>
T2	1.37 <sup>ab</sup>	2.66	48.49 <sup>ef</sup>	1.37 <sup>ab</sup>	2.67	48.64 <sup>e</sup>
T1	1.38 <sup>a</sup>	2.66	48.04 <sup>f</sup>	1.38 <sup>a</sup>	2.68	48.43 <sup>e</sup>
LSD (5 %)	0.014	0.019	0.696	0.014	0.012	0.606
Significance	**	**	**	**	**	**
CV (%)	0.62	0.42	0.82	0.62	0.26	0.72

Means followed by the same letter within a column are non-significantly different to each other at \* Significant at 5% level of significance ( $P < 0.05$ ) and \*\* highly Significant at 1% level of significance. ( $P < 0.01$ ) as established by LSD test; \* CM = Cattle manure in  $\text{t ha}^{-1}$ , RDF = Recommended Dose of inorganic NP Fertilizer,

Several studies have reported that CM plus mineral nutrient application resulted in reduced bulk density, higher soil organic carbon and hydraulic conductivity and improved soil structure and microbial communities (Andargachew,2014; Shiferaw,2014; Yadav *et al.*,2017; Kumar *et al.*,2017).

Bulk density normally decreases, as mineral soils become finer in texture. Therefore, the combined application of mineral NP and cattle manure decreased bulk density with increase in total pore space present in the soil and gives a good estimate of the porosity of the soil. Bulk density is of greater importance than particle density in understanding the physical behavior of the soil. Generally, soils with low bulk densities have favorable physical conditions. Similarly, Song (2016) reported that the decrease in bulk density obtained with manure treated soil was directly related to the increased organic matter, which played a significant role in reducing compaction of soil. The organic amendments promote the activity of soil fauna and play a major role in the buildup and stabilization of soil structure. In view of this, bulk density

values of the soils in the study areas were not too compact after crop harvest to limit root penetration and restrict movement of water and air.

Accordingly, the highest total porosity (51.78%) in Belg and (51.10%) in Meher was obtained from the application of 7.5 t ha<sup>-1</sup> CM in combination with 75% inorganic N and P and 50% inorganic N and P in which is statistically the same as application of 7.5 t ha<sup>-1</sup> CM in combination with 50% inorganic N and P in both Belg and Meher season. This increased porosity with the application of CM might be due to low soil compaction increase in organic matter leading to increased rapid multiplication of soil microorganisms, fastens the mineralization of plant nutrients in the soil. Similarly, Poeplau (2017) reported that soil with good organic matter content have higher porosity, and plays a critical role in establishing and maintaining soil fertility. On the other hand, the lowest total porosity of 48.04 % in Belg and 48.43 in Meher season were obtained from the control (Table 6).

**Soil Chemical Characteristics after Crop Harvest:** A comparison of soil fertility status prior to planting and after harvesting of potato shown that incorporation of cattle manure and mineral NP highly significantly ( $P < 0.01$ ) affects organic carbon content, total N, available P, soil pH, CEC, exchangeable acidity, exchangeable aluminum, exchangeable calcium, exchangeable magnesium, exchangeable potassium percent bases saturation, However, there was non-significant ( $P > 0.05$ ) effect on exchangeable sodium both in Belg and Meher seasons. (Table 4) indicating that there was an improvement of the fertility status of the soil due to an integrated nutrient management.

**Soil Ph:** Soil pH in pre-sowing soil was 5.01 in Belg and 4.8 in Meher season (Table 2) Soil pH value after crop harvest ranged from a minimum of 4.95 to a maximum of 5.15 in Belg and a minimum of 4.79 to a maximum of 4.94 in Meher season. The highest soil pH of 5.15 in Belg and 4.94 in Meher season after harvest was recorded at application 7.5 t ha<sup>-1</sup> CM, with 75% inorganic N and P. which is statistically at par with application of 7.5 t ha<sup>-1</sup> CM, with 50% inorganic N and P. and 7.5 t ha<sup>-1</sup> CM, with 25% inorganic N and P. which might be due to lime like materials such as calcium and magnesium present in manure increases soil pH. In line with this result, Suh *et al.* (2015) showed an increase in soil pH in plots applied with cattle manure following potato production.

Similarly, Ahmed *et al.* (2015) also found that using of animal manure such as cattle manure has positively beneficial effects on changes the soil pH where increases or decreases in the pH, depending on the type of soil and properties of organic fertilizer while the lowest soil pH (4.95) in Belg and (4.79) in Meher season was obtained at zero application of nutrients which is the same as application of 100% NP, 2.5 t ha<sup>-1</sup> CM 25% NP, 2.5 t ha<sup>-1</sup> CM +50% NP in Belg season and 100% NP, 2.5 t ha<sup>-1</sup> CM +25% NP in Meher season. non-amendment of cattle

manure and mineral NP removal of organic amendments can increase reduced pH, of soil this is because organic amendments can increase Al availability to plants by increasing the H<sup>+</sup> release of soils. The observed reduced following non-application of treatments with organic amendments (e.g. cattle manure) and inorganic is understood to reduce decomposition (less negative charge of adsorption sites) and complexation of soluble Al and Fe by organic molecules. Therefore, the lesser the organic and inorganic amendments the greater the decrease in Al adsorption capacity of soil this increase the release of more H<sup>+</sup> and decline in soil pH. similar results were also reported by Haynes and Mokolobate, (2001) reduction in the decomposition of organic residues causes a reduction in soil pH

Table 7. pH of soil after harvest of potato as influenced by the application of cattle manure and inorganic NP in Belg and Meher season at Masha district, south-western Ethiopia

Treatment*	Belg	Meher
	pH (1:2.5 H <sub>2</sub> O)	pH (1:2.5 H <sub>2</sub> O)
2.5 t ha <sup>-1</sup> CM+25% NP	4.98 <sup>fg</sup>	4.82 <sup>h</sup>
2.5 t ha <sup>-1</sup> CM+50% NP	4.99 <sup>efg</sup>	4.83 <sup>efg</sup>
2.5 t ha <sup>-1</sup> CM+75% NP	5.00 <sup>def</sup>	4.84 <sup>ef</sup>
5 t ha <sup>-1</sup> CM+25% NP	5.02 <sup>def</sup>	4.86 <sup>de</sup>
5 t ha <sup>-1</sup> CM+50% NP	5.03 <sup>bcde</sup>	4.87 <sup>cde</sup>
5 t ha <sup>-1</sup> CM+75%NP	5.04 <sup>bcd</sup>	4.89 <sup>bcd</sup>
7.5 t ha <sup>-1</sup> CM+25% NP	5.06 <sup>bc</sup>	4.90 <sup>abc</sup>
7.5 t ha <sup>-1</sup> CM+50% NP	5.07 <sup>b</sup>	4.90 <sup>abc</sup>
7.5 t ha <sup>-1</sup> CM+75% NP	5.15 <sup>a</sup>	4.94 <sup>a</sup>
100% NP	4.96 <sup>g</sup>	4.80 <sup>gh</sup>
Control	4.95 <sup>g</sup>	4.79 <sup>h</sup>
LSD (5 %)	0.544	0.037
Significance	**	**
CV (%)	0.046	0.453

Means followed by the same letter within a column are non-significantly different to each other at \* Significant at 5% level of significance ( $P < 0.05$ ) and \*\* highly Significant at 1% level of significance. ( $P < 0.01$ ) as established by LSD test; \* CM = Cattle manure in t ha<sup>-1</sup>

Organic carbon: Total organic carbon before planting was 1.2% in Belg and 1.02% in Meher (Table 2). After harvesting, it ranged from a minimum of 18.8 to a maximum of 21.1g kg<sup>-1</sup> in Belg season and from a minimum of 10.36 to a maximum of 12.72 g kg<sup>-1</sup> in Meher season (Table 8).

The highest values at 7.5 t ha<sup>-1</sup> CM +75% RDF in both Belg and Meher season followed by 7.5 t ha<sup>-1</sup> CM +50% RDF. After harvesting, application of 7.5 t ha<sup>-1</sup> CM with75% RDF improved the soil organic carbon by 12.23% and 22.78% as compared to zero application of treatments in Belg and Meher seasons respectively. indicating that these amendments positively affected the organic matter.

Increasing levels of cattle manure are linked to greater increased in soil organic matter due to enhanced soil microbial action increase soil moisture, soil structure, nutrient retention,

aeration, water holding capacity of amended soil This indicated that the integrated application of 7.5 t ha<sup>-1</sup> CM with 75% recommended rates of inorganic NP had the capacity of improving the soil carbon content, both in Belg and Meher seasons.

Kumar *et al.* (2017) reported that the combined application of CM from 12.5 to 25 t ha<sup>-1</sup> with (75% and 100%) recommended doses of NPK (150:60:100) fertilizers in potato has increased the soil carbon content from (5.4-5.8 g kg<sup>-1</sup>) and (5.3 to 5.7 g kg<sup>-1</sup>) after potato harvest. In addition, Kumar *et al.* (2012) described that application of 50 % recommended dose of NPK (60:60:30) through inorganic fertilizers and remaining 50% recommended dose of N (60 kg N ha<sup>-1</sup>) from cattle manure manures gave higher organic carbon pool (12.61 g kg<sup>-1</sup>) as compared to 100% recommended dose of NPK through fertilizers after three years of potato cultivation

Similarly, Wang *et al.* (2012) reported that combined use mineral fertilizers and organic amendments had greater soil organic carbon content than those receiving only mineral fertilizers in paddy soil. In addition, crop residue.

Total nitrogen: Total nitrogen in pre-sowing soil was 0.1% in Belg and 0.08% in Meher season (Table 2). soil total nitrogen after harvest ranged from a minimum of 1.24 to a maximum of 1.71g kg<sup>-1</sup>in Belg and a minimum 0.92 to a maximum of 1.51 g kg<sup>-1</sup> in Meher season from zero application of nutrients to the application of 7.5 t ha<sup>-1</sup> CM with 75% recommended rate of mineral NP which resulted in improvement of soil nitrogen after harvest by 37.45% and 63.85%over zero application of nutrients (Table 8). This shows cattle manure prevents loss of mineral nitrogen into the environment. by gradually releases essential nutrients over a period through regulating the rate of conversion of cattle manure to total nitrogen in addition, to release of nutrients into the soil after decomposition

This result was in concomitant with the finding of Lemanowicz *et al.* (2014) reported that increasing the application CM from zero to 20, 40, and 80 t ha<sup>-1</sup> and mineral nitrogen from 0, 40, 80 and 120 kg ha<sup>-1</sup> increased the total nitrogen content of soil as compared to the soil sampled from the treatments fertilized with the zero CM and mineral NP rate. In addition, Girma *et al.* (2017) reported that application of 14 t ha<sup>-1</sup> CM combines with 55kg ha<sup>-1</sup> and 20 t ha<sup>-1</sup> mineral NP fertilizers improve soil total nitrogen supply by 48%. Similarly, Biruk *et al.* (2015) reported that the treatment receiving CM at 30 t ha<sup>-1</sup> + N at 120 kg ha<sup>-1</sup> + 92 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup>. Improves total N by 67 % in the soil after harvest of potato as compared to zero application of mineral and cattle manure. Inlined with this Kumar *et al.* (2008) also stated that the total soil nitrogen content after harvest of potato increased in response to the application of CM and mineral fertilizer than 100% inorganic fertilization. from the present result, the increasing levels of cattle manure and mineral NP application increase the total nitrogen content of soil after harvest.

Table 8. Organic carbon ( $\text{g kg}^{-1}$ ), total nitrogen ( $\text{g kg}^{-1}$ ) and available phosphorus ( $\text{mg kg}^{-1}$ ) content of soil after potato harvest as influenced by the application of cattle manure and mineral NP in Belg and Meher seasons in Masha district, south-western Ethiopia.

Treatment*	Belg			Meher		
	Organic carbon ( $\text{g kg}^{-1}$ )	Total nitrogen ( $\text{g kg}^{-1}$ )	Available phosphorus ( $\text{mg kg}^{-1}$ )	Organic carbon ( $\text{g kg}^{-1}$ )	Total nitrogen ( $\text{g kg}^{-1}$ )	Available phosphorus ( $\text{mg kg}^{-1}$ )
2.5 t ha <sup>-1</sup> CM+25% NP	18.9 <sup>c</sup>	13.7 <sup>bcd</sup>	6.24 <sup>bcd</sup>	10.4 <sup>b</sup>	0.98 <sup>gf</sup>	5.55 <sup>cd</sup>
2.5 t ha <sup>-1</sup> CM+50% NP	19.1 <sup>c</sup>	13.7 <sup>bcd</sup>	6.24 <sup>bcd</sup>	10.7 <sup>b</sup>	1.06 <sup>ef</sup>	5.57 <sup>bcd</sup>
2.5 t ha <sup>-1</sup> CM+75% NP	19.2 <sup>c</sup>	13.7 <sup>bcd</sup>	6.24 <sup>bcd</sup>	10.7 <sup>b</sup>	1.12 <sup>de</sup>	5.64 <sup>bcd</sup>
5 t ha <sup>-1</sup> CM+25%NP	19.3 <sup>c</sup>	13.7 <sup>bc</sup>	6.24 <sup>bcd</sup>	10.94 <sup>b</sup>	1.15 <sup>cde</sup>	5.71 <sup>bcd</sup>
5 t ha <sup>-1</sup> CM+50% NP	19.3 <sup>c</sup>	14.8 <sup>bc</sup>	6.77 <sup>abc</sup>	10.96 <sup>b</sup>	1.164 <sup>cd</sup>	5.77 <sup>bc</sup>
5 t ha <sup>-1</sup> CM+75%NP	19.4 <sup>c</sup>	15.1 <sup>abc</sup>	6.95 <sup>ab</sup>	11.00 <sup>b</sup>	1.24 <sup>c</sup>	5.78 <sup>bc</sup>
7.5 t ha <sup>-1</sup> CM+25%NP	20 <sup>bc</sup>	15.4 <sup>ab</sup>	7.12 <sup>ab</sup>	11.13 <sup>b</sup>	1.38 <sup>b</sup>	5.82 <sup>bc</sup>
7.5 t ha <sup>-1</sup> CM+50% NP	20.8 <sup>ab</sup>	15.8 <sup>ab</sup>	7.30 <sup>a</sup>	12.33 <sup>a</sup>	1.46 <sup>ab</sup>	5.94 <sup>ab</sup>
7.5 t ha <sup>-1</sup> CM+75% NP	21.1 <sup>a</sup>	17.1 <sup>a</sup>	7.55 <sup>a</sup>	12.72 <sup>a</sup>	1.51 <sup>a</sup>	6.27 <sup>a</sup>
100% NP	18.83 <sup>cd</sup>	13.1 <sup>cd</sup>	5.88 <sup>cd</sup>	10.42 <sup>b</sup>	0.96 <sup>g</sup>	5.37 <sup>de</sup>
Control	18.78 <sup>d</sup>	12.4 <sup>d</sup>	5.53 <sup>d</sup>	10.36 <sup>b</sup>	0.92 <sup>g</sup>	5.003 <sup>e</sup>
LSD (5 %)	0.144	2.14	0.96	0.98	0.09	0.38
Significance	**	**	**	**	**	**
CV (%)	4.33	8.67	8.61	5.21	4.92	3.92

Available P: Available P in pre-sowing soil was 5.5  $\text{mg kg}^{-1}$  in Belg and 5  $\text{mg kg}^{-1}$  in Meher season (Table 2) while soil available P after harvest ranged from minimum of 5.53  $\text{mg kg}^{-1}$  to maximum of 7.55  $\text{mg kg}^{-1}$  in Belg and minimum of 5 to maximum of 6.27  $\text{mg kg}^{-1}$  in Meher seasons compared to the initial soil content of available P and the control (Table 8). The highest available P in Belg (7.55  $\text{mg kg}^{-1}$ ) and in Meher (6.27  $\text{mg kg}^{-1}$ ) season were obtained from the application of 7.5 t ha<sup>-1</sup>CM in combination with 50 and 75% mineral NP fertilizers, respectively

The increase in available P with increased level of CM and mineral NP application was probably related to the addition of various organic sources improve the soil physio-chemical properties that created the favorable condition in rhizosphere, also the secretion of certain enzymes and auxins and other growth-promoting substances and manure also helps in producing intermediate compounds that interact with phosphorus-fixing cations such as aluminum, iron, etc. Thereby reducing P adsorption capacity.

This result was in agreement with the investigation of Baviskar *et al.* (2011) and Balemi (2012) where they reported the increase in available P with the application of cattle manure which might have helped in releasing a higher amount of P from the soil. Similarly, Kesarwani (2017) reported that application of 5t ha<sup>-1</sup> CM with 120:60:40 NPK  $\text{kg ha}^{-1}$  enhanced P availability, increased overall treatments than sole use of mineral fertilizer as well. In addition, Shethet *al.*(2017) showed that integrated use of 15 t ha<sup>-1</sup> cattle manure combined with 100%

recommended dose of inorganic fertilizers (75: 50: 75 NPK kg ha<sup>-1</sup>) is efficient than application of inorganic fertilizers alone in improving growth and yield in sweet potato cultivation. The present result was also in agreement with the findings of Singh *et al.* (2017) reported that the increased available P content of soil with the application of cattle manure might be due to release of organic acids during decomposition which in turn helped in releasing phosphorus. Generally, the available P status of the soils in the control at both study seasons was very low, even below the critical level (10 ppm) (Isreal *et al.*, 2016) which indicates that the low soil P was among the factors highly limiting the productivity of the soils in the study area. This result was in agreement with the investigation of Song *et al.* (2017) where they reported that continuous use of organic manure and chemical fertilizer on soil improves the available phosphorus status of the soil.

Table 9. The combined effect of cattle manure and inorganic NP fertilizer on exchangeable acidity (cmol(+) kg<sup>-1</sup>soil) and exchangeable aluminum (cmol(+) kg<sup>-1</sup>soil) in Masha district in Belg and Meher seasons, south-western Ethiopia.

Treatment*	Belg		Meher	
	Exchangeable acidity	Exchangeable aluminum	Exchangeable acidity	Exchangeable aluminum
2.5 t ha <sup>-1</sup> CM+25% NP	3.12 <sup>c</sup>	2.19 <sup>ab</sup>	2.91 <sup>ab</sup>	1.98 <sup>ab</sup>
2.5 t ha <sup>-1</sup> CM+50% NP	2.87 <sup>d</sup>	2.16 <sup>ab</sup>	2.79 <sup>b</sup>	1.90 <sup>b</sup>
2.5t ha <sup>-1</sup> CM+75% NP	2.79 <sup>d</sup>	2.15 <sup>ab</sup>	2.52 <sup>c</sup>	1.71 <sup>c</sup>
5 t ha <sup>-1</sup> CM+25% NP	2.78 <sup>d</sup>	2.10 <sup>ab</sup>	2.44 <sup>cd</sup>	1.66 <sup>cd</sup>
5 t ha <sup>-1</sup> CM+50% NP	2.60 <sup>e</sup>	1.99 <sup>bc</sup>	2.34 <sup>cde</sup>	1.59 <sup>cde</sup>
5 t ha <sup>-1</sup> CM+75% NP	2.48 <sup>ef</sup>	1.81 <sup>c</sup>	2.25 <sup>def</sup>	1.53 <sup>def</sup>
7.5 t ha <sup>-1</sup> CM+25% NP	2.36 <sup>f</sup>	1.54 <sup>d</sup>	2.20 <sup>ef</sup>	1.50 <sup>ef</sup>
7.5 t ha <sup>-1</sup> CM+50% NP	2.34 <sup>f</sup>	1.52 <sup>d</sup>	2.17 <sup>ef</sup>	1.47 <sup>ef</sup>
7.5 t ha <sup>-1</sup> CM+75% NP	2.32 <sup>f</sup>	1.46 <sup>d</sup>	2.08 <sup>f</sup>	1.41 <sup>f</sup>
100% NP	3.41 <sup>b</sup>	2.21 <sup>a</sup>	3.10 <sup>a</sup>	2.11 <sup>a</sup>
Control	3.72 <sup>a</sup>	2.26 <sup>a</sup>	3.09 <sup>a</sup>	2.10 <sup>a</sup>
LSD (5 %)	0.173	0.196	0.227	0.154
Significance	**	**	**	**
CV (%)	3.63	5.93	5.25	5.25

Means followed by the same letter within a column are non-significantly different to each other at \* Significant at 5% level of significance ( $P < 0.05$ ) and \*\* highly Significant at 1% level of significance. ( $P < 0.01$ ) as established by LSD test; \* CM = Cattle manure in t ha<sup>-1</sup>.

On the other hand, the lowest exchangeable acidity of 2.32 cmol (+) kg<sup>-1</sup> soil in Belg and 2.08 cmol (+) kg<sup>-1</sup> soil in Meher season was recorded at application of 7.5 t ha<sup>-1</sup> cattle manure with 75% inorganic N and P which is statistically at par with application of 7.5 t ha<sup>-1</sup> cattle manure with 50% inorganic N and P and 7.5 t ha<sup>-1</sup> cattle manure with 25% inorganic N and P which might be that the addition of cattle manures, can increase perception of exchangeable Al which might have decrease exchangeable acidity and aluminum. Similarly Narambuye and Haynes (2012) reported that an increase in soil pH leads to reduction of soluble exchangeable Al which, in turn, decreases exchangeable acidity.

Exchangeable aluminum: The exchangeable aluminum pre-sowing soil was 2.01 cmol (+) kg<sup>-1</sup> soil in Belg 2.46 cmol (+) kg<sup>-1</sup> soil in Meher season (Table 2) while, soil total nitrogen after harvest ranged from 2.26 to 1.46 cmol (+) kg<sup>-1</sup> in Belg and to 2.10 to 1.41 cmol (+) kg<sup>-1</sup> in Meher season (Table 9)

The highest exchangeable aluminum in Belg (2.26 cmol (+) kg<sup>-1</sup> soil) and Meher (2.10 cmol (+) kg<sup>-1</sup> soil) was recorded at control which might be due to a low organic matter reduces the affinity for Al<sup>+3</sup> leads to increases the solubility and availability of aluminum. Similarly, Ayalew and Dejene (2012) indicated that treatments with no fertilizer application be it inorganic N and Por organic gave highest exchangeable acidity and Al while the lowest exchangeable Al and acidity was obtained at the combined application of 46 kg N + 40 kg P + 50 kg K + 20 t ha<sup>-1</sup> CM, However, the lowest exchangeable Al<sup>+3</sup> in Belg (1.465 cmol (+) kg<sup>-1</sup>) and Meher (1.475 cmol (+) kg<sup>-1</sup>) were recorded at combined application 7.5 t ha<sup>-1</sup> CM with 75% inorganic N and P 7.5 t ha<sup>-1</sup> CM with 50% inorganic N and P and 7.5 t ha<sup>-1</sup> CM with 25% inorganic N and P which could be due to decomposition of cattle manure may add more Ca and Mg that may increase the affinity of aluminum, might result in a reduction of soluble and exchangeable aluminum. This result is in agreement with the work of Wong and Swift (2003) reported decreased Al solubility because of the exchange between soil and organic matter with high cattle manure increases the soil pH this will decrease the solubility of Al

Similarly, Mccauley *et al.* (2017) stated that the reduction in exchangeable Ac and Al partly relates to the increase in soil pH because the increase in pH resulted in precipitation of exchangeable and soluble Al as insoluble Al hydroxides, thus reducing concentrations of Al in the soil solution

Effects of treatments on cation exchange capacity and exchangeable bases : Cation exchange capacity (CEC): Cation exchange capacity in pre-sowing soil was 20 cmol (+) kg<sup>-1</sup> in Belg and 19.3 cmol (+) kg<sup>-1</sup> in Meher. While (Table 2). Cation exchange capacity after harvest ranged from 16.35 cmol (+) kg<sup>-1</sup> to 22.04 cmol (+) kg<sup>-1</sup> in Belg and from 19.09 cmol (+) kg<sup>-1</sup> to 23.96 cmol (+) kg<sup>-1</sup> in Meher season (Table 9). Accordingly, the highest CEC of 22.04 cmol (+) kg<sup>-1</sup> in Belg and 23.96 cmol (+) kg<sup>-1</sup> in Meher season were obtained from the combined application of 7.5 t ha<sup>-1</sup> CM with 75% inorganic N and P and 7.5 t ha<sup>-1</sup> CM with 50% inorganic N and P fertilizers in Belg and 7.5 t ha<sup>-1</sup> CM with 75% inorganic N and P, 7.5 t ha<sup>-1</sup> CM with 50% inorganic N and P and 7.5 t ha<sup>-1</sup> CM with 25% inorganic N and P fertilizers in Meher season (Table 9). On the other hand, the lowest CEC observed in the control plots which could be attributed to the leaching of bases from the soil due to the high rainfall of the study areas, which were not replenished since there was no fertilizer provided to these plots thus decrease in organic matter content reduces site for carbon storage and sequestration, lowered microbial biomass in the soil and reduced nutrient reserves. The application of 7.5 t ha<sup>-1</sup> CM in combination with 75% minral NP both in Belg and Meher could be due to the contribution of organic matter from CM to the high organic matter content in the plots than the other treatments. in line with this result, Tolanur (2002) reported that cation exchange capacity significantly increased with increase in organic manure in conjunction with inorganic fertilizer the mineral fertilizers which are readily taken up by the crop is subsisted by the use of organic amendments such as cattle manure has been recognized generally as an effective means for

improving soil aggregation, structure and fertility causes increasing microbial diversity and populations, results in increasing surface charges, thus in turn leads to improvement in cation exchange capacity of soil.

Similarly, Dorneles *et al.* (2015) reported the highest CEC from the non-tillage system because of high organic matter as compared to another tillage system. Cation exchange capacity (CEC) of the soil is one of the most important components of soil fertility evaluation to supply nutrient cations to the soil solution and possible response to fertilizer application (Sonon *et al.*, 2014). This indicates basic cations existing in a given soil and the natural and/or anthropogenic activities, acting upon these cations, thereby influencing the CEC of the soil. (Xavier, 2015).

Table 10. Effect of application of inorganic NP and cattle manure on CEC (cmol (+) kg<sup>-1</sup> soil, Ca<sup>+2</sup>(cmol (+) kg<sup>-1</sup> soil, Mg<sup>+2</sup> (cmol (+) kg<sup>-1</sup>soil) of soil after crop harvest in Belg and Meher season in Masha district, south-western Ethiopia

Treatment *	Belg			Meher		
	CEC	Ca <sup>+2</sup>	Mg <sup>+2</sup>	CEC	Ca <sup>+2</sup>	Mg <sup>+2</sup>
2.5 t ha <sup>-1</sup> CM+25% NP	17.41 <sup>efg</sup>	6.39 <sup>efg</sup>	2.76 <sup>bcde</sup>	21.33 <sup>bcde</sup>	7.19 <sup>bcde</sup>	1.40 <sup>efg</sup>
2.5 t ha <sup>-1</sup> CM+50% NP	17.01 <sup>efg</sup>	6.22 <sup>efg</sup>	2.70 <sup>cde</sup>	20.88 <sup>cde</sup>	6.98 <sup>cde</sup>	1.34 <sup>fg</sup>
2.5 t ha <sup>-1</sup> CM+75% NP	17.98 <sup>def</sup>	6.63 <sup>def</sup>	2.76 <sup>bcde</sup>	21.35 <sup>bcde</sup>	7.18 <sup>bcde</sup>	1.48 <sup>def</sup>
5 t ha <sup>-1</sup> CM+25% NP	18.36 <sup>de</sup>	6.79 <sup>de</sup>	2.89 <sup>abcd</sup>	22.20 <sup>abcd</sup>	7.54 <sup>abcd</sup>	1.54 <sup>de</sup>
5 t ha <sup>-1</sup> CM+50% NP	18.86 <sup>cd</sup>	7.00 <sup>cd</sup>	2.98 <sup>abc</sup>	22.84 <sup>abc</sup>	7.80 <sup>abc</sup>	1.61 <sup>cd</sup>
5 t ha <sup>-1</sup> CM+75% NP	19.28 <sup>cd</sup>	7.18 <sup>cd</sup>	2.99 <sup>abc</sup>	22.96 <sup>abc</sup>	7.86 <sup>abc</sup>	1.67 <sup>cd</sup>
7.5 t ha <sup>-1</sup> CM+25% NP	20.01 <sup>bc</sup>	7.49 <sup>bc</sup>	3.07 <sup>ab</sup>	23.49 <sup>ab</sup>	8.09 <sup>ab</sup>	1.77 <sup>bc</sup>
7.5 t ha <sup>-1</sup> CM+50% NP	20.88 <sup>a</sup>	7.86 <sup>ab</sup>	3.13 <sup>a</sup>	23.91 <sup>a</sup>	8.26 <sup>a</sup>	1.90 <sup>ab</sup>
7.5 t ha <sup>-1</sup> CM+75% NP	22.04 <sup>a</sup>	8.36 <sup>a</sup>	3.14 <sup>a</sup>	23.96 <sup>a</sup>	8.28 <sup>a</sup>	2.06 <sup>a</sup>
100% NP	16.68 <sup>fg</sup>	6.08 <sup>fg</sup>	2.59 <sup>de</sup>	20.16 <sup>de</sup>	6.67 <sup>cde</sup>	1.23 <sup>fg</sup>
Control	16.35 <sup>fg</sup>	5.95 <sup>g</sup>	2.44 <sup>e</sup>	19.09 <sup>e</sup>	6.22 <sup>de</sup>	1.27 <sup>g</sup>
LSD (5 %)	1.36	0.57	0.33	2.31	0.98	0.19
Significance	**	**	**	**	**	**
CV (%)	4.29	4.86	6.78	6.17	7.71	7.09

Means followed by the same letter within a column are non-significantly different to each other at \* Significant at 5% level of significance ( $P < 0.05$ ) and \*\* highly Significant at 1% level of significance. ( $P < 0.01$ ) as established by LSD test; \* CM = Cattle manure in t ha<sup>-1</sup>

Exchangeable calcium: Exchangeable calcium in pre-sowing soil was 6.5 cmol (+) kg<sup>-1</sup> soil in Belg 6.3 cmol (+) kg<sup>-1</sup> soil in Meher season (Table 2) while, soil exchangeable calcium after harvest ranged from 5.95 to 8.36 cmol (+) kg<sup>-1</sup> in Belg and from 6.22 to 8.28 cmol (+) kg<sup>-1</sup> in Meher season (Table 10). The highest exchangeable Ca<sup>+2</sup> (8.36 cmol (+) kg<sup>-1</sup> soil) in Belg and (8.28 cmol (+) kg<sup>-1</sup> soil) in Meher season was recorded in the combined application, of a high dose of cattle manure (7.5 t ha<sup>-1</sup>) with 75% inorganic N and P fertilizers that showed an increase of 40.36% and 33% in Belg and Meher season, respectively, as compared to zero application of nutrients. The elevated calcium levels were associated with decreasing exchangeable aluminum in this soil. with a dramatic increase in pH causes improvement in Exchangeable Calcium in conformity with this result, Shivan and Tolanur (2002) reported that application of cattle manure in combination with inorganic fertilizers improved significantly



exchangeable calcium when organic materials combine together with inorganic materials. On the other hand the lowest Exchangeable calcium ( $5.95 \text{ cmol (+) kg}^{-1}$  soil) in Belg and ( $6.22 \text{ cmol (+) kg}^{-1}$ ) in Meher was obtained at zero application of nutrients which is statistically the same as application of 100% NP,  $2.5 \text{ t ha}^{-1}$  CM + 25% NP,  $2.5 \text{ t ha}^{-1}$  CM+50% NP in Belg season and 100% NP,  $2.5 \text{ t ha}^{-1}$  CM+25% NP,  $2.5 \text{ t ha}^{-1}$  CM + 50% NP and  $2.5 \text{ t ha}^{-1}$  CM+75% NP in Meher season

Exchangeable magnesium: Exchangeable magnesium in pre-sowing soil was  $2.1 \text{ cmol (+) kg}^{-1}$  in Belg  $1.4 \text{ cmol (+) kg}^{-1}$  in Meher season while soil Exchangeable magnesium after harvest ranged from 2.44 to 3.14 in Belg and 1.27 to 2.07 in Meher season (Table 2). The highest exchangeable magnesium content of soil after harvest ( $3.14 \text{ cmol (+) kg}^{-1}$  soil) in Belg and ( $2.07 \text{ cmol (+) kg}^{-1}$  soil) in Meher were recorded at combined application of cattle manure with 75% inorganic N and P. (Table 9) This may be due to amending soil with cattle manure raises soil organic matter which increases the soil pH leads to greater activities of soil bacteria, which in turn improves availability of soil nutrients and the application of magnesium from any external sources improves the soil exchangeable magnesium. This result is partly in line with work of Daniel and Nigussie (2006) reported that amending the soil with cow dung improves the soil exchangeable magnesium content of soil after harvest. In contrast, the lowest exchangeable magnesium content of the soil after harvest ( $2.44 \text{ cmol (+) kg}^{-1}$  soil) in Belg and ( $1.275 \text{ g cmol (+) kg}^{-1}$  soil) in Meher were obtained at the zero application of nutrients.

Exchangeable potassium: Exchangeable potassium in pre-sowing soil was  $0.42 \text{ cmol (+) kg}^{-1}$  soil in Belg  $0.36 \text{ cmol (+) kg}^{-1}$  soil in Meher (Table 2) while, soil Exchangeable potassium after harvest ranged from  $0.31 \text{ cmol (+) kg}^{-1}$  soil to  $0.43 \text{ cmol (+) kg}^{-1}$  soil in Belg and to  $0.26$  to  $0.34 \text{ cmol (+) kg}^{-1}$  in Meher season. The highest exchangeable potassium of  $0.43 \text{ cmol (+) kg}^{-1}$  soil in Belg and  $0.34 \text{ cmol (+) kg}^{-1}$  soil in Meher was obtained at the combined application of  $7.5 \text{ t ha}^{-1}$  CM, with 75% inorganic N ( $165 \text{ kg N ha}^{-1}$ ) and P ( $137 \text{ kg P}_2\text{O}_5 \text{ ha}^{-1}$ ). (Table 11). Which is statistically the same as  $7.5 \text{ t ha}^{-1}$  CM with 50% recommended dose of inorganic N ( $165 \text{ kg N ha}^{-1}$ ) and P ( $137 \text{ kg P}_2\text{O}_5 \text{ ha}^{-1}$ ) and  $7.5 \text{ t ha}^{-1}$  CM with 25% recommended dose of inorganic N ( $165 \text{ kg N ha}^{-1}$ ) and P ( $137 \text{ kg P}_2\text{O}_5 \text{ ha}^{-1}$ ) while the lowest exchangeable potassium in both seasons was recorded at zero rates organic and inorganic fertilizer (Table 10). The higher status of exchangeable potassium both in Belg and Meher seasons of higher dosage plots indicates amendment of soil with external fertilizers increases exchangeable K with an increase in nutrient dosage similarly Abay and Sheleme (2012) reported that periodic assessments for potassium status of the soil and the effect of the crop to potassium are essential and as potato is a highly K demanding crop,

Table 11. Exchangeable K<sup>+</sup>, Na<sup>+</sup> and Pbs of soil after crop harvest as influenced by the combined application of cattle manure and inorganic NP in Belg and Meher seasons, in Masha district, south-western Ethiopia

Treatment*	Belg			Meher		
	Exchangeable K <sup>+</sup> cmol (+) kg <sup>-1</sup> soil	Exchangeable Na <sup>+</sup> cmol (+) kg <sup>-1</sup> soil	Percent base saturation (%)	Exchangeable K <sup>+</sup> cmol (+) kg <sup>-1</sup> soil	Exchangeable Na <sup>+</sup> cmol (+) kg <sup>-1</sup> soil	Percent base saturation (%)
2.5 t ha <sup>-1</sup> CM+25% NP	0.36 <sup>de</sup>	0.02	46.73 <sup>ghi</sup>	0.28 <sup>cde</sup>	0.20	48.73 <sup>bcd</sup>
2.5 t ha <sup>-1</sup> CM+50% NP	0.36 <sup>de</sup>	0.02	47.03 <sup>fgh</sup>	0.29 <sup>bcde</sup>	0.21	48.95 <sup>abc</sup>
2.5 t ha <sup>-1</sup> CM+75% NP	0.37 <sup>cd</sup>	0.03	47.40 <sup>efg</sup>	0.29 <sup>bcde</sup>	0.21	48.95 <sup>abc</sup>
5 t ha <sup>-1</sup> CM+25% NP	0.38 <sup>bcd</sup>	0.03	47.65 <sup>def</sup>	0.31 <sup>abcd</sup>	0.22	49.34 <sup>abc</sup>
5 t ha <sup>-1</sup> CM+50% NP	0.38 <sup>bcd</sup>	0.04	47.95 <sup>cde</sup>	0.30 <sup>abc</sup>	0.23	49.60 <sup>ab</sup>
5 t ha <sup>-1</sup> CM+75% NP	0.39 <sup>bd</sup>	0.04	48.20 <sup>cd</sup>	0.32 <sup>abc</sup>	0.23	49.65 <sup>ab</sup>
7.5 t ha <sup>-1</sup> CM+25% NP	0.40 <sup>abc</sup>	0.05	48.57 <sup>cd</sup>	0.33 <sup>ab</sup>	0.23	49.84 <sup>ab</sup>
7.5 t ha <sup>-1</sup> CM+50% NP	0.42 <sup>ab</sup>	0.06	49.01 <sup>ab</sup>	0.33 <sup>a</sup>	0.24	49.99 <sup>a</sup>
7.5 t ha <sup>-1</sup> CM+75% NP	0.43 <sup>a</sup>	0.07	49.54 <sup>a</sup>	0.34 <sup>a</sup>	0.24	50.08 <sup>a</sup>
100%NP	0.35 <sup>de</sup>	0.01	46.48 <sup>hi</sup>	0.28 <sup>de</sup>	0.20	48.33 <sup>cd</sup>
Control	0.31 <sup>e</sup>	0.01	46.20 <sup>i</sup>	0.26 <sup>e</sup>	0.19	47.65 <sup>d</sup>
LSD (5 %)	0.04	0.01	0.73	0.035		1.13
Significance	**	Ns	**	**	Ns	*
CV (%)	6.17	23.12	0.90	6.78	7.04	1.35

Means followed by the same letter within a column are non-significantly different to each other at \* Significant at 5% level of significance ( $P < 0.05$ ) and \*\* highly Significant at 1% level of significance. ( $P < 0.01$ ) as established by LSD test; \* CM = Cattle manure in t ha<sup>-1</sup>.

Percent bases saturation: percent bases saturation: percent bases saturation in pre-sowing soil was 45.4% in Belg and 41.7% in Meher (Table 2) while, soil percent bases saturation after harvest ranged from 46.202%to 49.54% in Belg and to 47.65% to 50.08 % in Meher season. The highest percent bases saturation of 49.54(%) in Belg and 50.08(%) in Meherseasons) was obtained at the combined application of 7.5 t ha<sup>-1</sup> CMwith 75% and 50% (RDF) and 50.08(%) was obtained at 7.5 t ha<sup>-1</sup> CM, with 75% (RDF) which is statically the same as all rates except Control, 100% NP,2.5 t ha<sup>-1</sup> CM+25% NP, 2.5 t ha<sup>-1</sup> CM+50% While the lowest percent bases saturation in both seasons was recorded at zero rates of either organic or inorganic fertilizer which is statistically the same as application of 100% NP and 2.5 t ha<sup>-1</sup> CM+25% NP in Belg and Meher seasons. (Table 11).

The higher status percent bases saturation both in Belg and Meher seasons may be due to the addition of organic and inorganic amendment together increases organic matter, surface area for exchange site and this causes favorable environment for plant growth by reducing leaching, enhancing growth in density and length of root hairs Similarly Wasihun *et al.* (2015) reported that treatments with high organic matter have high PBS.

As conclusion, the soil test results after potato harvest revealed significant increase in soil, pH, EC, TN, available P, CEC, exchangeable calcium, exchangeable magnesium, exchangeable potassium, porosity and tuber yield of potato, on the other hand, significant in reducing soil bulk and particle density, exchangeable acidity and Al when CM was applied in combination with inorganic NP than the use of either zero application of NP or 100% recommended rate of NP at both seasons. Among the treatments, the combined application of 7.5 t CM with 75% inorganic NP fertilizer was superior. Hence, the use of CM (7.5 t ha<sup>-1</sup>) with 75% recommended rate of N (123.75 kg N ha<sup>-1</sup>) and P (103.05 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup>) was found to be appropriate to improve the physico-chemical properties of the soil and to increase the productivity of potato in the study areas as compared to the 100% recommended rate of NP (165 kg N ha<sup>-1</sup> and 137 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup>) alone.

#### ACKNOWLEDGMENTS

We thank Ministry of Education (MoE) of Ethiopia, for the financial support. Teppi and Hawassa Soil Research Center for timely analyzing the soil and cattle manure samples and material support the sustainable land management (SLM) project. In Masha, District Sheka zone is highly acknowledged for providing Cattle manure and sponsoring the laboratory cost incurred for this work.

#### REFERENCES

- Abay Ayalew and Sheleme Beyene. 2012. Characterization of soils and response of potato (*Solanum tuberosum* L.) to the application of potassium at Angacha in southern Ethiopia. *Journal of Biochemistry and Bioinformatics*, 2(2): 46–57.
- Adejumo, S.A., Togun, A.O. Adediran J.A. and M.B. Ogundiran, 2010. Effects of compost application on remediation and the growth of maize planted on lead-contaminated soil, Department of Crop Protection and Environmental Biology, University of Ibadan, Ibadan. Nigeria.
- Agegnehu Getachew and Amede Tilahun 2017. Integrated Soil Fertility and Plant Nutrient Management in Tropical Agro-Ecosystems: *A Review*. *Pedosphere*, 27(4): 662–680.
- Ahmed, A. A., Zaki, M. F., Shafeek, M. R., Helmy, Y. I and El-bakery, M. M. H. A. 2015. Integrated Use of Cattle Manure and Mineral Nitrogen Fertilizer on Growth, Yield, and Quality of Potato (*Solanum tuberosum* L.). *International Journal of Current Microbiol. Applied. and Science*, 4(10): 325–349.

- Balemi, T. 2012. Effect of integrated use of cattle manure and inorganic fertilizers on tuber yield of potato in Ethiopia. *Journal of Soil Science and Plant Nutrition*,12(2): 253-261
- Banti Gizaw. 2009. Geographical location and agro-ecological variation of Kaffa Zone. MSc. Thesis. Addis Ababa University, Addis Ababa, Ethiopia.
- Baruah, T.C. and Barthakulh, H.P. 1997. *A textbook of soil analysis*. Vikas Publishing House. New Delhi, India. 334p.
- Bayu, W., and Gebeyehu, S. 2013. Effect of Cattle Manure and Inorganic Fertilizers on the Growth, Yield and Moisture Stress Tolerance of Rain-fed Lowland Rice. *American Journal of Research Communication*, 1(4): 275–301.
- Biruk-Masrie, Nigussie-Dechassa, Tamado Tana, Yibekal Alemayehu and Bekele Abebi. 2015. Effect of Application of Cattle Manure and Mineral NP Fertilizers on Yield, and Nutrient Uptake of Potato in North-Eastern Ethiopia. *Journal of Science and Sustainable Development*, 3(1): 1-23.
- Black, C.A. 1965. Methods of soil analysis. *American Society of Agronomy*,1: 1570-1572.
- Bray, R.H. and Kurtz, 1945. Determination of total, organic and available forms of phosphorus in soils. *Soil Science*, 59: 39–45.
- Chapman, H.D., 1965. Cation exchange capacity. In: C.A. Black, L.E. Ensminger and F.E.Clark (Eds). *Methods of soil analysis*. Agronomy. 9: 891-901. Am. Soc. Agro., Inc., Madison, Wisconsin.
- Compaore E., Cattan P and Taonda JB. 2011. Effect of Continuous Mineral and Organic Fertilizer Inputs and Plowing on Groundnut Yield and Soil Fertility in a Groundnut–Sorghum Rotation in Central Burkina Faso. In: Bationo A., Waswa B., Okeyo J., Maina F., Kihara J. (eds) *Innovations as Key to the Green Revolution in Africa*. Springer, Dordrecht
- CSA (Central Statistical Authority). 2016. Agricultural sample survey 2015/16. Volume I Report on area and production of major crops for private peasant holdings, meher season. Statistical bulletin 578, Central Statistical Agency, Addis Ababa, Ethiopia.
- Daniel Mekonnen and Nigussie Dechassa. 2006. Effects of integrated nutrient management on the agronomic performance of potato (*Solanum tuberosum* L.) and fertility of Nitosol at Bako. MSc. Thesis. Haramaya University, Haramaya, Ethiopia

- Day, P.R.1965. Hydrometer method of particle size analysis. In: C.A. Black (ed.). Methods of soil analysis. Agronomy Part I, No. 9. American Society of Agronomy. Madison, Wisconsin, USA. pp.562-563.
- Dejene M. and Lemlem M. 2012. Integrated agronomic crop management to improve teff productivity under terminal drought. pp. 235- 254. In: I. Md. M. Rahman and H. Hasegawa, (eds.), Water Stress, In Technology Open Science
- Dell'Angelo, J., D'Odorico, P., and Rulli, M. C. 2017. Threats to sustainable development posed by land and water grabbing. Current Opinion in Environmental Sustainability. 26-27: 120-128
- Efthimiadou, A., Bilalis, D., Karkanis, and Froud, B. 2010. Combined organic and inorganic fertilization enhances soil quality and increased yield, photosynthesis, and sustainability of sweet maize crop. Australian Journal of Crop Science, 4 (9): 722-729.
- EthioSIS (Ethiopia Soil Information System). 2014. Soil fertility status and fertilizer recommendation atlas for Tigray regional state, Ethiopia. July 2014, Addis Ababa, Ethiopia
- FAO (Food and Agriculture Organization). 2006. Plant nutrition for food security: A guide for integrated nutrient management. FAO, fertilizer and plant nutrition bulletin 16, Rome. Italy.
- Fekadu Asfaw. 2016. Effect of Integrated Soil Amendment Practices on Growth and Seed Tuber Yield of Potato (*Solanum tuberosum* L.) at Jimma Arjo, Western Ethiopia. *Journal of Natural Sciences Research*, 6:1-26.
- Gafar, A.F., Yassin, M., Ibrahim, D. and Yagoob, S.O. 2012. Effect of different (bio-organic and mineral) fertilizers on some yield components of rice. Universal Journal of Agricultural Research, 2(2): 67-70.
- Getachew Angachew. 2009. Ameliorating effects of organic and mineral fertilizers on crop productivity and soil properties on reddish-brown soils. pp. 127-150. In: Proceeding of the tenth Conference of the Ethiopian Society of Soil Science. 25-27 March 2009, held at EIAR, Addis Ababa, Ethiopia.
- Haynes, R. J., and Mokolobate, M. S. 2001. Amelioration of Al toxicity and P deficiency in acid soils by additions of organic residues: a critical review of the phenomenon and the mechanisms involved. *Nutrient Cycling in Agroecosystems*, 59: 47-63.

- Hazelton, P. and Murphy, B., 2007. Interpreting Soil Test Results: What do all the numbers Mean 2<sup>nd</sup> (eds.). CSIRO Publishing. 152p.
- Hillel, D. 2004. Introduction to Environmental Soil Physics. Elsevier Academic Press.
- Israel Zewide, Ali Mohammed and Solomon Tulu Tadesse. 2012. Effect of Different Rates of Nitrogen and Phosphorus on Yield and Yield Components of Potato (*Solanum tuberosum* L.) at Masha District, South-western Ethiopia. *International Journal of Soil Science*, 7(4): 146-156.
- Jackson, M.L. 1958. Soil Chemical Analysis. Prentice-Hall, Inc., Engle Wood Cliffs. New Jersey. pp. 183 - 204.
- Jones, J.B. 2003. Agronomic Handbook: Management of Crops, Soils, and their fertility. CRC Press LLC, Boca Raton, FL, USA. 482p.
- Kabira, J. N. 2017. Effect of inorganic fertilizer and cattle manure on growth and yield of two Kenyan potato varieties, *International Journal of Agronomy and Agricultural Research*, 10(1): 65-72.
- Kang, Y., Hao, Y., Shen, M., Zhao, Q., Li, Q., and Hu, J. 2016. Impacts of supplementing chemical fertilizers with organic fertilizers manufactured using pig manure as a substrate on the spread of tetracycline resistance genes in soil. *Ecotoxicology and Environmental Safety*, 130: 279-288.
- Kesarwani, A. 2017. Effect of Integrated Nutrient Management on Growth and Yield Parameters of Maize (*Zea mays* L) As well as Soil Physicochemical Properties. *Biomedical Journal of Scientific and Technical Research*, 1(2): 2-7.
- Kuldeep Balai, M Jajoria, R Verma, P Deewan and SK Bairwa. 2016. Nutrient content, uptake, quality of chickpea and fertility status of soil as influenced by fertilization of Phosphorus and Zinc. *Journal of Pharmacognosy and Phytochemistry*, 6(1): 392-398.
- Kumar Manoj, Baishaya L.K, Ghosh D.C, Gupta V.K, Dubey S.K, Das Anup and Patel D.P. 2012. Productivity and soil health of potato (*Solanum tuberosum* L.) field as influenced by organic manures, inorganic fertilizers, and biofertilizers under high altitudes of Eastern Himalayas. *Journal of Agricultural Science*, 4: 223-234.
- Kumar Praveen, Pandey SK, Singh SV, Singh BP, Rawal S and Kumar Dinesh. 2008. Evaluation of nutrient management options for potato processing cultivars. *Potato Journal*, 35 (1-2): 46-52.

- Kumar, P., Kumar, A., Kumar, N., Ahamad, A., and Verma, M. K. (2017). Effect of Integrated Nutrient Management on Productivity and Nutrients Availability of Potato. *International Journal of Current Microbiology and Applied Sciences*. 6(3), 1429–1436.
- Landon, J.R. 1991. Booker Tropical Soil Manual: A Handbook for Soil Survey and Agricultural Land Evaluation in the Tropics and Subtropics. Longman Scientific and Technical, Essex, New York. 474p.
- Lesschen, J.P., Stoorvogel, J.J., Smaling, E.M.A., Heuvelink, G.B.M. and Veldkamp, A. 2007. A spatially explicit methodology to quantify soil nutrient balances and their uncertainties at the national level. *Nutrient Cycling in Agroecosystems*, 78: 111-131.
- Lindsay, W. L., and Norvell, W. A. 1978. Development of a DTPA Soil Test for Zinc, Iron, Manganese, and Copper. *Soil Science Society of American Journal*, 42(3):421–428.
- Mahajan, A., Bhagat, R.M. and Gupta, R.D. 2008. Integrated nutrient management in sustainable rice-wheat cropping system for food security in India. *Journal of Agriculture*, 6 (2): 29-32.
- Mccauley, A., Jones, C., and Olson-Rutz, K. 2017. Soil pH and Organic Matter. *Nutrient Management*, Module No 16.
- MoARD (Ministry of Agriculture and Rural Development). 2012. Crop Development Department of Crop Variety Register. Issue No. 10. Addis Ababa, Ethiopia.
- Najm, A. A., Hadi, M. R. H. S., Fazeli, F., Darzi, M. T., and Rahi, A. 2012. Effect of Integrated Management of Nitrogen Fertilizer and Cattle Manure on the Leaf Chlorophyll, Yield, and Tuber Glycoalkaloids of Agraria Potato. *Communications in Soil Science and Plant Analysis*, 43(6): 912–923.
- Narambuye F. X and Haynes. R. J 2006. Effect of organic amendments on soil pH and AL solubility and use of laboratory indices to predict their liming effect. *Soil Science*, 171(10): 754–763.
- Ndegwa PM and Thompson SA.2001. Integrating composting and vermicomposting in the treatment and bioconversion of solids. *Bioresource Technol* 76: 07–112.
- Okalebo JR, Guthua KW and Woomer PJ 2002. Laboratory methods of soil and plant analysis a working manual. Tsbf-ciat and sacred Africa, Nairobi

- Opala, P. A., Okalebo, J. R and Othieno, C. O. 2012. Effects of Organic and Mineral Materials on Soil Acidity and Phosphorus Availability in a Soil Incubation Study. *International Scholarly Research Network Agronomy*, 1: 1-10.
- Page, A.L. 1982. *Methods of Soil Analysis*. Part II. Chemical and Microbiological Properties. Madison, USA.
- Paul, J., Asstt, A., Choudhary, A. K., Sharma, S., Savita. S and Bohra, M .2016. Potato production through bio-resources: Long-term effects on tuber productivity, quality, carbon sequestration and soil health in the temperate Himalayas. *Scientia Horticulture*, 213: 152–163.
- Poeplau, C., Vos, C., and Don, A. 2017. Soil organic carbon stocks are systematically overestimated by misuse of the parameters of bulk density and rock fragment content. *Soil*. 3(1): 61–66.
- SAS (Statistical Analysis System). 2003. SAS Version 9. 1.2 © 2002-2003". SAS Institute, Inc., Cary, North Carolina, USA.
- Saviozzi and Cardelli, 2014. Organic matter characteristics, biochemical activity and antioxidant capacity of soil amended with different organic materials. *Archives of Agronomy and Soil Science*, 60(1):119-131.
- Sheth, S. G., Desai, K. D., Patil, S. J., and Navya, K., L. 2017. Effect of integrated nutrient management on growth, yield, and quality of sweet potato [*Ipomoea batatas*( L .) Lam ], *International Journal of Chemical Studies*, 5(4): 346-349
- Shiferaw Boke. 2014. Effect of organic and inorganic fertilizer application and seedbed preparation on potato yield and soil properties on Alisols of Chencha. *International Journal of Natural Sciences Research*, 2(8): 123–132.
- Shubhadip D, Arindam S, Krishna Chaitanya<sup>1</sup>, A, Aritra S, Arup D and Riasen M. 2017. The response of Potato Crop to Integrated Nutrient Management in the Indo-Gangetic Alluvial Soils of, of West Bengal, India. *Journal of Experimental Agriculture International*, 16(3): 1–10.
- Song Ke, Yong Xue, Xianqing Zheng, Weiguang Lv, Hongxia Qiao, Qin n, and Jianjun Yang. 2017. Effects of the continuous use of organic manure and chemical fertilizer on soil mineral Phosphorus fractions in calcareous soil. *Scientific Reports*, 7: 1164.



- Song, K., Yang, J., Xue, Y., Lv, W., Zheng, X., and Pan, J. 2016. Influence of tillage practices and straw incorporation on soil aggregates, organic carbon, and crop yields in a rice-wheat rotation system. *Scientific Reports*, 6: 1-12.
- Sonon, L. S., Kissel, D. E., and Saha, U. 2014. Cation Exchange Capacity and Base Saturation. UGA Extension Circular, 1040:1-4.
- Suh, C., Meka, S. S., Ngome, A. F., Neba, D. A., Kemngwa, I. T., Sonkouat, A. D. and 1Njualem, D. 2015. Effects of organic and inorganic fertilizers on growth and yield of potato (*Solanum tuberosum* L.) In the western highlands of Cameroon. *International Journal of Development Research*, 5: 3584–3588.
- Suresh Kumar Kakraliya<sup>1</sup> R.D., Sandeep Kumar<sup>1</sup>, K.K. Choudhary, Jai Prakash and L.K. Singh. 2017. Integrated Nutrient Management for Improving, Fertilizer Use Efficiency, Soil Biodiversity and Productivity of Wheat in Irrigated Rice-Wheat Cropping System in Indo-Gangatic Plains of India *International Journal Current Microbiol. Applied Science*, 6(3): 152-163.
- SZARDO (Sheka Zone Agricultural and Rural Development Office) 2015. annual report
- Tekalign Tadesse, Haque, L. and Aduayi, E.A. 1991. Soil, Plant, Water, Fertilizer, Animal Manure and Compost Analysis Manual: Plant Science Division Working Document NO. 13. International Livestock Center for Africa. Addis Ababa, Ethiopia.
- Temesgen Gashaw; Amare Bantider and Hagos G/Silassie. 2014. Land Degradation in Ethiopia: Causes, Impacts and Rehabilitation Techniques. *Journal of Environment and Earth Science*, 4(9): 98–105.
- EPA (Environmental Protection Authority). 2012. National Report of Ethiopia, the United Nations Conference on Sustainable Development (Rio+20). Federal Democratic Republic of Ethiopia, Addis Ababa.
- Tolanur, S.I. 2002. Integrated Nutrient Management on physical conditions, soil fertility and productivity of crops in dryland Vertisol. *Journal of Karnataka Agricultural Science*, 14(3): 186- 187.
- Walkley, A. J. and Black, I. A. 1934. Estimation of soil organic carbon by the chromic acid titration method. *Soil Science*, 37: 29 - 38.
- Wang, S., Liang, X., Luo, Q., Fan, F and Chen, Y., Li, Z. 2012. Fertilization increases paddy soil organic carbon density. *Journal of Zhejiang University*, 13(4): 274–282.

- Wasihun Mengiste, Mukhtar Mohammed, and Teshome Yitebarek 2015. Evaluation of the effect of land use types on selected soil physico-chemical properties in Itang-key area of Gambella regional state of Ethiopia MSc. Thesis Haramaya University, Haramaya, Ethiopia.
- Wong, M. T. F., and Swift, R. S. 2003. Role of organic matter in alleviating soil acidity. *Handbook of Soil Acidity*. Marcel Dekker. New York. The USA.
- Xavier López-Medellín. 2015. Natural resource management to improve rural livelihoods in a natural protected area in Central Mexico: Exploring communities' attitudes on conservation *Journal of Soil and Water Conservation*, 14(2): 89-99.
- Yadav, R. K., Purakayastha, T. J., Khan, M. A., and Kaushik, S. C. 2017. The science of the Total Environment Long-term impact of manuring and fertilization on enrichment, stability, and quality of organic carbon in Inceptisol under two potato-based cropping systems. *The science of the Total Environment*, 609: 1535–1543.
- Yadav, S. K., Babu, S., Yadav, M. K., Singh, K., Yadav, G. S., and Pal, S. 2013. A Review of Organic Farming for Sustainable Agriculture in Northern India. *International Journal of Agronomy*, 1: 1–8.
- Zelege, G., Agegnehu, G., Abera, D., Rashid, S. 2010. Fertilizer and soil fertility potential in Ethiopia: Constraints and opportunities for enhancing the system. IFPRI, Addis Ababa, Ethiopia, 63p.