

Productive and physiological responses of West African dwarf goats fed diets containing gmelia leaf meal.

Respuestas productivas y fisiológicas de cabras enanas de África occidental alimentadas con dietas que contienen harina de hojas de gmelia.

Jiwuba, P.C.*, Onwujiariri, E.B. and Akazue, R.C.

Department of Animal Production Technology, Federal College of Agriculture, P.M.B.7008, Ishiagu, Ebonyi State, Nigeria.

*Corresponding author: jiwubapc@gmail.com

ABSTRACT

West African Dwarf (WAD) goat plays an important socio-economic role, hence the effect of 82 days of feeding Gmelina leaf meal (GLM) diets on feed intake, body weight change and blood indices of WAD goats were investigated. 36 WAD goats averaging 9.04kg in weight were randomly divided into four groups of nine goats each with three goats constituting a replicate. The groups were randomly assigned four experimental diets (T₁, T₂, T₃, and T₄) in a completely randomized design. Results indicated significant ($p < 0.05$) improvement for body weights and feed conversion ratio for T₂, T₃, and T₄. There was significant difference ($P < 0.05$) for all the haematological parameters except packed cell volume (PCV). White blood cells (WBC) was improved ($p < 0.05$) with incremental levels of GLM in the diets. All the serum

biochemical indices showed significant ($p < 0.05$) difference with glucose and cholesterol concentration reduced ($p < 0.05$) with incremental levels of GLM. It could be concluded that GLM can be included in the diets of goats up to 36% without deleterious effects on the performance of WAD goats.

Keywords: blood chemistry, body weight changes, feed intake, haematology, leaf meal, phytochemical compounds and WAD goats.

RESUMEN

La cabra enana de África occidental (WAD) juega un papel socioeconómico importante, por lo que se investigó el efecto de 82 días de alimentación con dietas de harina de hojas de Gmelina (GLM) sobre la ingesta de alimento, el cambio de peso corporal y los índices sanguíneos de las cabras WAD. Se dividieron al azar 36 cabras WAD con un peso promedio de 9,04 kg en cuatro grupos de nueve cabras cada uno con tres cabras que constituían una réplica. A los grupos se les asignaron al azar cuatro dietas experimentales (T1, T2, T3 y T4) en un diseño completamente aleatorizado. Los resultados indicaron una mejora significativa ($p < 0.05$) para el peso corporal y la tasa de conversión alimenticia para T2, T3 y T4. Hubo una diferencia significativa ($P < 0,05$) para todos los parámetros hematológicos excepto el volumen de células empaquetadas (PCV). Los glóbulos blancos (WBC) mejoraron ($p < 0.05$) con niveles incrementales de GLM en las dietas. Todos los índices bioquímicos séricos mostraron una diferencia significativa ($p < 0,05$) con la concentración de glucosa y colesterol reducida ($p < 0,05$) con los niveles incrementales de GLM. Se podría concluir que GLM puede incluirse en las dietas de cabras hasta en un 36% sin efectos deletéreos sobre el rendimiento de las cabras WAD.

Palabras clave: química sanguínea, cambios en el peso corporal, ingesta de alimento, hematología, harina de hojas, compuestos fitogénicos y cabras WAD.

INTRODUCTION

West African dwarf (WAD) goat is the most prolific ruminant in the south eastern Nigeria capable of increasing the animal protein intake. They are multipurpose animal, producing meat, milk, skin and manure. Their reputation for short gestation interval, early maturity, easy management, remarkable capacity to convert roughages into meat and milk, unique ability to adapt and maintain itself in harsh environments, low cost of production, ability trek long distances in search of feed, high dressing percentage, ability to adapt to a wide range of climatic conditions, high meat quality and their small body sizes make them indispensable livestock in Nigeria and beyond. The meat is generally accepted and consumed in Nigeria since there is no religious taboo against it. The demand for the meat is high and usually command higher market price than beef and pork (Jiwuba *et al.*, 2018a).

However, the production of this animal, with great potentials to alleviate animal protein inadequacy, is faced with various challenges. Jiwuba *et al.* (2017a) attributed high cost of feed and inadequate nutrient intake to undermine this goat breed in expressing their full potential. This may be as a result of little knowledge about the potentials of some non-conventional feed resources, the presence in some cases of anti-nutritional element and the lack of proper storage capabilities especially over long period (Ben Salem *et al.*, 2009). In line with this, interests have been shifted to the search for cheaper, available and nutritionally viable feedstuffs with some phyto-genic properties to enhance livestock production. Some leaf meals of tropical origin have been reported (Jiwuba *et al.*, 2017a; Jiwuba, 2018; Jiwuba *et al.*, 2018b) to yield relatively higher levels of crude protein, dry matter and lower crude fibre levels than most tropical forages.

Gmelina arborea is an evergreen perennial fast growing medicinal tree, of which are used as laxative and anthelmintic, improve appetite, useful in hallucination, piles, abdominal pains, burning sensations, fevers, and urinary discharge (Kumaresan *et al.*, 2014). The leaves are unconventional materials that can be explored for the production of feedstuff. The leaf is

one of the tree leaves considered as an important source of nutrient for ruminants and non-ruminants, especially in those areas with a pronounced dry season. Previous records by Nkwocha *et al.* (2014) and Offor (2014) on the nutrient profile of *Gmelina arborea* have shown that the leaf meals contained 18.00 - 20.05% crude protein, 14.40 - 15.05% crude fibre, ash 4.55% and fat 0.79%. Metabolizable energy values were found to be appreciable (1368 Kcal Kg⁻¹) an indication of its suitability as an energy source for livestock diets (Amata, 2014). The leaves, flower, roots and bark are used in medicine. They are useful in the treatment of blood diseases; thus the use of *Gmelina* leaf meal in the present study fits in the strategy of improving the productive and physiological responses of WAD goats. Therefore, this experiment was carried out to investigate the effects of graded levels of *Gmelina* leaf meal on the feed intake, body weight changes, haematological and serum biochemical indices of WAD goats.

MATERIALS AND METHODS

The research work was carried out at Sheep and Goat unit of Federal College of Agriculture Ishiagu, Ivo Local Government Area of Ebonyi State, Nigeria. The College is located at about three Kilometer (3km) away from Ishiagu main town. The College is situated at latitude 5° 6'N and longitude 7° 31'E with average rainfall of 1000-1600mm and a prevailing air temperature condition of 27-28°C relative humidity of about 88% respectively.

The *Gmelina* (*Gmelina arborea*) leaves were sourced within the College environs. Fresh, succulent, greenish non-over matured leaves were harvested to ensure lower value of lignin and higher nutrient availability. The leaves were air dried in batches to about 10% moisture before passing through a 10 mm hammer mill and further used in the formulation of the experimental diets. Four diets T₁, T₂, T₃, and T₄, were formulated. The *Gmelina* leaf

meal (GLM) was included at the levels of 0%, 12%, 24% and 36% for T₁, T₂, T₃, and T₄, respectively as presented in Table 1.

Thirty six (36) WAD goats of about 10 – 12 months of age and averaging 9.04 kg in weight were selected from the College herd for this experiment. The goats were randomly divided into four groups of nine animals each with three goats constituting a replicate. The groups were randomly assigned the four experimental diets (T₁, T₂, T₃, and T₄) in a completely randomized design (CRD). The animals were housed individually in a well-ventilated cement floored pens equipped with feeders and drinkers. Each animal received a designated treatment diet in the morning (8am) for 82 days. Feed offered was based on 3% body weight per day; the animals in addition were fed 1kg wilted chopped *Pennisetum purpureum* later in the day (5pm). Regular access to fresh drinking water was made available in accordance with the permission and stipulated guidelines of the Federal College of Agriculture, Ishiagu (FCAI) Animal Ethics Committee. Initial live weights of the animals were taken at the beginning of the feeding trial and weekly thereafter. Final live weight was obtained by weighing the goats at the end of the experiment. Daily weight gain, daily feed intake and feed conversion ratio were calculated.

Ten ml of blood samples were drawn from each animal on the last day of the study. The goats were bled through the jugular vein. The samples were separated into two lots and used for haematological and biochemical determinations. An initial 5ml was collected from each sample in labelled sterile universal bottle containing 1.0 mg/ml ethyldiamine tetracetic acid (EDTA) and used for haematological analysis. Another 5ml was collected over anti-coagulant free bottle and used for the serum biochemical studies. Serum biochemistry and haematological parameters were measured using Beckman Coulter Ac-T10 Laboratory Haematology Blood Analyzer and Bayer DCA 2000+ HbA1c analyzer, respectively. Mean corpuscular haemoglobin (MCH), mean corpuscular volume (MCV) and mean corpuscular haemoglobin concentrations (MCHC) were calculated.

All the sample of feed and test ingredients were analyzed for their proximate composition using the method of AOAC (2000). Gross energy was calculated using the formula $T = 5.72Z_1 + 9.50Z_2 + 4.79Z_3 + 4.03Z_4 \pm 0.9\%$; where T = Gross energy, Z_1 = Crude protein, Z_2 = Crude fat, Z_3 = Crude fibre, Z_4 = Nitrogen free extract (Nehring & Haelein 1973).

The results were analyzed using the Statistical Package for Social Sciences Window 17.0. One -way analysis of variance (ANOVA) was employed to determine the means and standard error. Treatment means were separated using Duncan's new multiple range test (Duncan, 1955).

RESULTS AND DISCUSSION

The proximate composition of the test diets and Gmelina leaf meal are presented in Table 1. The experimental diets and GLM were analyzed for dry matter (DM), crude protein (CP), crude fibre (CF), ash, ether extract (EE) and nitrogen free extract (NFE). EE and gross energy (GE) were significantly ($p < 0.05$) influenced by the treatments why DM, CP, CF, NFE and ash were not ($p > 0.05$) influenced. The DM of the experimental diets failed to follow a particular trend. The DM ranged between 91.62 and 92.36% with T_1 having the highest value (92.36%) and T_2 the lowest (91.62%). The CP increased progressively with incremental levels of GLM with the highest value of 14.64% recorded for T_4 with a corresponding lowest value of 13.04% recorded for T_1 . The CF and ash maintained a particular trend like the CP increasing with an increasing level of GLM. EE decreased ($p < 0.05$) progressively with incremental levels of GLM. NFE and GE values failed to follow a specific trend, increasing or decreasing with increasing levels of GLM.

The body weight changes and feed intake of West African dwarf goats fed diets containing Gmelina leaf meal is presented in Table 2. There were significant ($p < 0.05$) difference for final body weight, total weight gain (TWG), average daily weight gain (ADWG) and feed conversion ratio (FCR) among the experimental goats, while total feed intake (TFI)

and average daily feed intake (ADFI) remained similar ($p > 0.05$). There was improvement ($p < 0.05$) in the body weight among the treatment groups (T_2 , T_3 and T_4) in comparison with the control (T_1).

Table 1: Composition of the experimental diets for West African Dwarf Goats

Ingredient	Dietary levels				SEM	GLM	PKM
	T ₁	T ₂	T ₃	T ₄			
Gmelina leaf meal	0.00	12.00	24.00	36.00			
Cassava peel meal	45.00	45.00	45.00	45.00			
Palm kernel meal	45.00	33.00	21.00	9.00			
Ground nut meal	5.00	5.00	5.00	5.00			
Molasses	2.00	2.00	2.00	2.00			
Bone meal	2.00	2.00	2.00	2.00			
Common salt	1.00	1.00	1.00	1.00			
Total	100	100	100	100			
Analyzed							
Dry matter	92.36	91.62	91.73	91.89	17.27	93.90	91.81
Crude protein	13.04	13.43	13.75	14.64	11.47	18.32	16.78
Crude fibre	16.67	16.58	17.01	17.65	9.59	15.18	13.42
Ash	4.49	4.54	4.67	4.69	0.05	3.96	3.87
Ether Extract	4.88 ^a	3.67 ^a	3.06 ^a	2.33 ^b	0.02	1.16	6.12
NFE	53.28	53.40	53.24	52.58	12.11	55.28	50.62
GE (MJ/g)	4.15 ^a	4.06 ^{ab}	4.05 ^{ab}	4.02 ^b	0.04	4.11	4.22

^{a-b} means within a row with different superscript differ significantly ($p < 0.05$); SEM = Standard error of mean; GLM- Gmelina leaf meal; PKM = Palm kernel meal; NFE = Nitrogen free extract; GE = Gross energy

Table 2: Body weight changes and feed intake of WAD goats fed diets containing *Gmelina* leaf meal

Parameters	Treatments				SEM
	T ₁ (0%)	T ₂ (12%)	T ₃ (24%)	T ₄ (36%)	
Initial body weight (kg)	9.05	8.67	9.25	9.19	2.43
Final body weight (kg)	13.70 ^c	15.80 ^b	15.65 ^b	17.85 ^a	4.44
Total weight gain (kg)	4.65 ^c	7.13 ^b	6.40 ^b	8.66 ^a	1.01
Average Daily weight gain (g/day)	56.71 ^d	86.95 ^b	78.05 ^c	105.61 ^a	32.31
Total feed intake (kg)	28.74	30.41	29.31	28.95	11.73
Average Daily feed intake (g/day)	350.49	370.85	357.44	353.05	21.71
Feed conversion ratio	6.18 ^a	4.27 ^{bc}	4.58 ^b	3.34 ^c	0.06

^{a-c} means within a row with different superscript differ significantly ($p < 0.05$)

Table 3 shows the haematological indices of WAD goats fed Gmelina leaf meal containing diets. The WBC obtained in this study ($8.35 - 11.70 \times 10^9/l$) followed a particular trend, increasing with increasing levels of GLM. T₄ was significantly higher and better than T₃, T₂ and T₁, while T₂ and T₁ showed ($p > 0.05$) similarities. PCV failed to follow a particular trend, but was numerically highest in T₃, and lowest in T₁. The PCV values were not significantly ($P > 0.05$) affected across the treatment groups. Hb and RBC followed a similar trend as the WBC, increasing with increasing levels of GLM, with T₄ (9.35g/dl) and ($10.50 \times 10^{12}/l$) respectively having the highest value and T₁ (8.10 g/dl) and ($9.05 \times 10^{12}/l$) respectively the lowest. The MCH tended to decrease with increasing levels of GLM, with T₁ (7.00 pg) having the highest value and T₄ (5.95 pg) the lowest. MCHC was similar ($p > 0.05$) for T₁ and T₂ then differed ($p < 0.05$) from T₃ and T₄. T₄ tended to be superior and higher than other treatment groups. MCV, decreased with increasing levels of GLM, with T₁ (24.40 fl) having the highest value and differed ($p < 0.05$) from T₃ and T₄, with T₄ (19.55 fl) having a corresponding lowest value.

Table 3: The effect of Gmelina leaf meal on Haematology of West African Dwarf goats

Parameters	Treatment levels				SEM
	T ₁ (0%)	T ₂ (12%)	T ₃ (24%)	T ₄ (36%)	
White blood cell (x10 ⁹ /1)	8.35 ^c	8.90 ^c	10.70 ^b	11.70 ^a	1.51
Packed cell volume (%)	30.90	31.40	31.60	31.40	11.16
Haemoglobin (g/dl)	8.10 ^c	8.20 ^c	8.70 ^b	9.35 ^a	4.19
Red blood cell (x10 ¹² /1)	9.05 ^b	9.16 ^b	10.10 ^a	10.50 ^a	2.23
MCH (pg)	7.00 ^a	7.75 ^a	6.15 ^b	5.95 ^b	1.77
MCHC (%)	25.65 ^c	26.65 ^c	27.35 ^b	29.40 ^a	5.75
MCV (fl)	24.40 ^a	24.25 ^a	20.50 ^b	19.55 ^c	8.80

^{a-c} Means within a row with different superscript differed significantly (P<0.05); MCH = Mean Corpuscular Haemoglobin; MCHC= Mean Corpuscular Haemoglobin Concentration; MCV = Mean Corpuscular Volume; SEM = Standard Error of Means.

The serum biochemical indices of WAD goats fed Gmelina leaf meal are shown in Table 4. The total protein was significantly (P<0.05) higher for the treatment groups (T₂, T₃ and T₄) in comparison with the control group (T₁). The albumen showed significant (P<0.05) difference across the treatment groups with T₁ having the lowest value and T₄ with the highest value. Glucose had a progressive decrease across the treatments with incremental levels of GLM in the diets. Glucose also showed (P>0.05) similarities between T₁ and T₂ while T₃ and T₄ differed significantly (P<0.05) different across the treatments. Urea concentration had a linear decrease (p<0.05) from T₁ to T₄. The serum creatinine values of goats on diet T₁ and T₂ were significantly higher (p < 0.05) to those on diets T₃ and T₄. Cholesterol values for the control diet were lower (p< 0.05) to those on diet T₂, T₃ and T₄.

Table 4: Serum biochemistry of West African Dwarf goat fed Gmelina leaf meal

Parameters	Treatments				SEM
	T ₁ (0%)	T ₂ (12%)	T ₃ (24%)	T ₄ (36%)	

Total protein (g/dl)	6.46 ^d	6.94 ^c	7.35 ^b	8.13 ^a	0.23
Albumin (g/dl)	2.70 ^d	2.79 ^c	3.06 ^b	3.33 ^a	0.18
Glucose (mg/dl)	74.50 ^a	74.00 ^a	65.50 ^b	61.50 ^c	2.68
Urea (mg/dl)	17.06 ^a	15.88 ^b	15.12 ^c	13.16 ^d	0.53
Creatinine (mg/dl)	1.71 ^a	1.69 ^a	1.65 ^b	1.63 ^b	0.01
Cholesterol (mg/dl)	96.41 ^a	93.05 ^b	85.18 ^c	84.64 ^c	1.91

^{a-d} Means within a row with superscript differ significantly (P<0.05).

The dry matter range of 91.62 – 92.36% recorded in this study compared well with the range of 91.32-91.74% and 89.95-91.44% reported by Jiwuba *et al.* (2018a) for WAD goats fed yellow root cassava peel-centrosema leaf meal and Jiwuba *et al.* (2018c) for WAD goats fed cassava root sievate - cassava leaf meal based diets respectively. The crude protein and gross energy values in this study falls within the requirement for goats as recommended by ARC (1980), NRC (1981), and Norton (1994). The crude fibre range of 16.67 – 17.65 % obtained in this study followed a particular trend, increasing with increasing levels of GLM in the diets. The values however, compared with 16.83-18.63% for WAD goats fed *Moringa oleifera* leaf meal in their diets as reported by Jiwuba *et al.* (2017a). Adequate supply of dietary fibre tends to increase chyme chewing, salivation, reduces digestive problems, and promotes intestinal motility (Jiwuba, 2018) in ruminants. The ash values of 4.49 - 4.69% followed a particular pattern increasing with increasing levels of GLM; hence indicating that high mineral profile of GLM. The decreasing (p<0.05) values obtained in this study for EE and GE with incremental levels of GLM may be attributed to higher EE and GE values reported for PKM. The results of the proximate analysis of the GLM favourably compared with the reports of Abdu *et al.* (2012) and Jiwuba *et al.* (2016a) for the same leaf meal. The proximate

composition of PKM obtained in this study compared well with the reported values of Ezieshi and Olumo (2007) and Boateng et al. (2008).

The significantly high ADWG value at 36% inclusion level could be attributed to enhanced utilization of the diets; a view corroborated by Ukanwoko & Okehilem (2016) who reported improved body weight gain and feed conversion ratio in WAD goats fed on Gmelina leaf meal. The improved weight gain may also be attributed to the high nutrient profile of GLM. Okagbare *et al.* (2004) earlier reported GLM as a rich source of nutrients. This may have contributed to improving the nutritional quality of the diets which may have supported the enhanced performance of the goats. The non-significant high feed intakes observed among the treatment groups may indicate high palatability of the diets. The superior FCR of diet T₄ over the other diets is a reflection of the observed higher growth rate and higher feed utilization of the goats fed the respective diet.

The improvement of the white blood cells (WBC) among the T₂, T₃ and T₄ groups in comparison with the T₁ group may be attributed to the medicinal and pharmacological properties of GLM; a view corroborated by Kumaresan *et al.* (2014) and Deepthi *et al.* (2015). Furthermore, the WBC range of 8.35 to 11.70 x10¹²/L. fell within the range of 4 to 13 x10¹²/L as reported by Fraser & Mays (1986) for apparently healthy goat. This indicated that the animals were healthy; hence decreased WBC count below the normal range (leukocytopenia) is an indication of allergic conditions and certain parasitism, while elevated values (leukocytosis) indicate the existence of a recent infection (Jiwuba *et al.*, 2016b). Hb concentration obtained in this study, generally appeared to support high oxygen carrying capacity in the blood of the goats since they fell within the physiological range of 8-12 g/dl reported by Fraser & Mays (1986). The RBC counts (9.05-10.50 x 10¹²/l) reported in this study fell within the range of 8-18 x10¹²/l reported by Fraser & Mays (1986), 7- 15 g/dl reported by Daramola *et al.* (2005), 9.98 – 10.88 g/dl reported by Jiwuba *et al.* (2017a). The improved RBC counts recorded for goats in the T₄ diets gave clear evidence of non-

vulnerability to anaemia-related diseases by the goats. The MCH and MCV values obtained in this study fell within the normal physiological range of 5.2-8.0 (Pg) and 16-25 (fl) as reported by Fraser & Mays (1986) and Jiwuba *et al.* (2016c) respectively for apparently healthy goats. The reduced MCH counts recorded for goats in the T₃ and T₄ diets present a likely susceptibility to hypochromic anaemia-related disease conditions by these goats; since MCH is the average mass of haemoglobin per red blood cell in a blood sample. Ryan (2010) however opined that MCV has to be considered along with the MCH since MCV affects the content of haemoglobin present per cell. This thus indicated that the T₄ diets could present a likely susceptibility to anaemia-related disease conditions by the goats if fed for a longer time. The values of MCHC indicated that the nutritional quality of the ration was not compromised. Also the MCHC in this study fell within normal physiological range of 30 – 36 g/dl reported by Fraser & Mays (1986); hence a clear indication of absence of anemia among the experimental goats.

The total protein in this study fell within the range of 6.3-8.5g/dl reported by Daramola *et al.* (2005), for apparently healthy WAD goats. The serum protein obtained in this study is however comparable with the reported values of 7.3 g/dl and 6.2-7.3 g/dl by Taiwo & Ogunsanmi (2003) and Ukanwoko (2016) respectively for WAD goats. The higher serum protein observed in T₄ may be attributed to the higher dietary protein of the diet as evidenced in the chemical composition of the diets. This is in agreement with the findings of Jiwuba *et al.* (2016d) who observed a positive relationship between serum protein and dietary protein. This further indicated better utilization of the dietary proteins by the T₄ goats. The range of 2.70-3.33 g/dl for serum albumen obtained in this study were within the normal physiological of 2.7–3.9 g/dl reported by Kaneko *et al.* (2008) and favourably compared with 2.75-3.2 g/dl reported for WAD goats by Ukanwoko (2016). According Jiwuba *et al.* (2017b) albumin functions chiefly in the regulation of colloidal osmotic pressure of the blood, assist in the movement of fatty acids, hormones, bilirubin, cations and drugs in the blood. Nevertheless, the normal albumin concentration reported in this study gave evidence of dietary protein quality and adequacy among the treatment animals. Serum glucose concentration is of great

importance in assessing the efficiency of sugar absorption in animals. The level of concentration of glucose in the goats were lower among the T₄ goat, which may indicate high glucagon and low insulin concentrations in the blood. However, the range of 61.50 -74.50 mg/dl reported in this study is within the referral range of 50-75mg/dl reported by Kaneko *et al.* (2008) for goat; thus an indication of absence of high glucagon and low insulin concentration in the body of the experimental goat. The range of urea concentration (13.16-17.06mg/dl) recorded in this study compared well with 13.11-17.85 mg/dl reported by Jiwuba *et al.* (2016c) for WAD goats. The recorded values also fell within the reference range of 10-20mg/dl reported by Kaneko *et al.* (2008) for goats. This suggested that the kidneys and liver of the goats were well functioning. However, the high serum urea nitrogen recorded in T₁ may be attributed to increased catabolism of amino acids, when proteins of lower biological values are fed to animals. This thus affirms earlier reports (Jiwuba *et al.*, 2017a; Jiwuba *et al.*, 2018b) that leaf meal produces protein of higher quality. The creatinine values (1.63-1.71mg/dl) recorded in this study were higher than 0.48-0.88mg/dl reported by Jiwuba *et al.* (2017a) for WAD goats, but however, fell within the baseline serum creatinine range of 1.0-1.8mg/dl reported by Kaneko *et al.* (2008) for goats. The within normal physiological range obtained in this study implies that the animals have proper functioning kidneys. The blood cholesterol level in this study ranged from 84.64-96.41mg/dl with T₄ showing lower cholesterol level as against the T₁ group with the highest cholesterol level respectively. The serum cholesterol concentration recorded in this study were higher than those reported by Jiwuba *et al.* (2016c) for WAD but fell within the referral serum cholesterol concentration of 80/130 mg/dl for apparently healthy goats as reported by Kaneko *et al.* (2008). This is an indication that the goats were healthy and not prone to heart related disease conditions, since a high level of serum cholesterol is an indicator of cardio-vascular related diseases.

It is concluded from this study that Gmelina leaf meal can be included in the diets of goats up to 36% without deleterious effects on the growth performance, haematological and serum biochemical parameters of West African Dwarf goats.

REFERENCES

- Abdu SB, Hassan MR, Jokthan GE, Adamu HY, Yashim SM, Yusuf K, 2012. Effect of varied inclusion levels of Gmelina (*Gmelina arborea*) leaf meal on intake, digestibility and nitrogen in Red Sokoto bucks fed on sorghum glum based complete diets. *Advances in Agriculture, Sciences and Engineering Research* 2 (2): 79 – 84.
- Amata IA, 2014. The use of non-conventional feed resources (NCFR) for livestock feeding in the tropics: a review. *Journal of Global Biosciences* 3: 604-613.
- AOAC, 2000. Animal feed. In *AOAC Official Methods of Analysis (6th Edition)* 5 -15. AOAC International, Washington DC USA.
- ARC, 1980. The nutrient requirement of ruminant livestock. CABI Farnham Royal U.K.
- Ben Salem HAH, Niprilo A, Nefzauoi A, 2009. Polyethylene glycol in concentrate or feed blocks to deactivate condensed tannis in *Acacia Cyanophylla* Lindl forage: Effects on intake digestion and growth by barbina lambs. *Animal science* 75: 127 - 135.
- Boateng M, Okai, DB, Baah J Donkoh A, 2008. Palm kernel cake extraction and utilisation in pig and poultry diets in Ghana. *Livestock Reseach for Rural Development* 20(7).
- Daramola JO, Adelofo AA, Fatiba IA, Soladoye AO, 2005. Haematological and biochemical parameters of West African Dwarf goats. *Live. Res. for Rural Development* 17 (8): 20-25
- Deepthi P, Harini A, Prakash LH, 2015. A Review on Gambhari (*Gmelina Arborea* Roxb.). *Journal of Pharmacognosy and Phytochemistry* 4: 127 – 132.
- Duncan DB, 1955. Multiple Ranges and multiple F-tests. *Biometrics* 11: 1 – 42.

- Ezieshi EV, Olomu JM, 2007. Nutritional evaluation of palm kernel meal types: 1. Proximate composition and metabolizable energy Values. African Journal of Biotechnology 6 (21); 2484-2486.
- Fraser CM, Mays A, 1986. The Merck Veterinary Manual. A handbook of diagnosis, therapy and disease prevention and control for the Veterinarian. Sixth Edition. Merck & Co., Inc. Rahway, New Jersey, USA, 905–908.
- Jiwuba PC, 2018. Effect of pawpaw (*Carica papaya*) leaf meal on productive parameters of growing rabbits. Agricultural Science and Technology 10 (2):102 – 106.
- Jiwuba PC, Ahamefule FO, Ogbuewu IP, Ikwunze K, 2017a. Blood chemistry and haematology of West African Dwarf goats fed *Moringa oleifera* leaf meal (MOLM) in their diets. Comparative Clinical Pathology 26 (3): 621-624.
- Jiwuba PC, Ezenwaka LC, Ikwunze K, Nsidinanya NO, 2016c. Blood profile of West African Dwarf goats fed provitamin A cassava peel-centrosema leaf meal based diets. Analele Stiintifice ale Universitatii,, Alexandru Ioan Cuz”, Sectiunea Genetica si Biologie Moleculara TOM XVII, Fascicula. 3:27-134.
- Jiwuba PC, Ikwunze K, Dauda E, Ugwu DO, 2016b. Haematological and serum biochemical indices of growing Rabbits fed diets containing varying levels of *Moringa oleifera* leaf meal. British Biotechnology Journal 15(2): 1-7.
- Jiwuba PC, Jiwuba LC, Onyekwere MU, 2017b. Growth performance, haematology and serum biochemistry of West African dwarf sheep fed cassava peel - oil palm leaf meal based diets in a hot humid tropics. Agricultura 14(1-2): 37-44. DOI: 10.1515/agricultura-2017-0016
- Jiwuba PC, Ogbuewu IP, MNwachukwuguru K, 2018b. Performance and economy of production of broilers fed Siam weed (*Chromolaena odorata*) leaf meal (SWLM). Tropical Animal Health and Production.

- Jiwuba PC, Onunka BN, Nweke JC, 2018c. Influence of supplemental cassava root sieviate - cassava leaf meal based diets on carcass and economics of production of West African dwarf goats. Sustainability, Agri, Food and Environmental Research 6(4): 40-54.
- Jiwuba PC, Onwujiariri EB, Kadurumba OE, 2018a. Carcass yield, organ response and cost/benefit evaluation of West African dwarf goats fed yellow root Cassava Peel-Centrosema leaf meal based diets. Nig. J. Anim. Prod. 45(2): 342 – 351.
- Jiwuba PC, Ugwu DO, Kadurumba OE, Dauda E, 2016a. Haematological and serum biochemical indices of weaner rabbits fed varying levels of dried *Gmelina arborea* leaf meal. International Blood Research and Reviews 6: 1-8.
- Kaneko JJ, Harvey JW, Bruss ML, 2008. Clinical Biochemistry of Domestic Animals. 6th Ed., Academic Press.
- Kumaresan P, Jeyanthi KA, Kalaivani R, 2014. Biochemical evaluation of anti diabetic activity of aqueous extract of *Gmelina arborea* in Alloxan induced albino rats. International Journal of Herbal Medicine 2: 90-94
- Nehring K, Haelien GWF, 1973. Feed evaluation and calculation based on net energy. J. Anim.Sci, 36 (5): 949 – 963.
- Nkwocha GA, Agbabiaka LA, Beketin TO, Anukam KU, 2014. Effect of graded levels of *Gmelina arborea* leaf meal on the performance of grower Rabbits. International Journal of Agricultural Science 4: 370-373.
- Norton BWB, Lowry C, Sweeney MC, 1994. The nutritive value of *Leucaena* specie. Paper presented at Int'l. Workshop on *Leucaena*. R.D. Bogor, Indonesia. 20-29.
- NRC, 1981. Nutrient Requirements for goats: Angora, dairy and meat goat in temperate and tropical countries. National Research Council; National Academy of Science Press, Washington DC.

Offor CE, 2014. Phytochemical and proximate analyses of dry *Gmelina arborea* leaves. International journal of current research and academic review 2: 101-105

Okagbare GO, Akpodiete OJ, Esiekpe O, Onagbesan OM, 2004. Evaluation of *Gmelina arborea* leaves supplemented with grass) *Panicum maximum* and *Pennisetum purpureum*) as feed for West African dwarf goats. Trop. Anim. Health Pro. 36: 593-598.

Ryan DH, 2010. Examination of blood cell. Lichtman MA, Kipps TJ, Seligsohn U et al. eds Williams Hematology. 8th ed. New York: the McGraw-Hill companies, Inc.

Taiwo VO, Ogunsanmi AO, 2003. Haematology, plasma, whole blood and erythrocyte biochemical values of clinically healthy captive-reared grey duiker (*Sylvicapra grimmia*) and West African dwarf sheep and goats in Ibadan, Nigeria. Israel Journal of Veterinary Medicine 58(2-3):

Ukanwoko AI, 2016. Blood profile of West African dwarf (WAD) goats fed *Gmelina* (*GmelinaArborea*) leaf meal based diets. Int'l Journal of Agric. and Rural Dev. 19(1): 2500-2505.

Ukanwoko AI, Okehilem OV, 2016. Effect of *Gmelina* (*Gmelina arborea*) leaf meal based diets on growth performance of West African dwarf bucks. Asian Journal of Animal Sciences.

Received: 11th November 2020; Accepted: 01th July 2021; First distribution: 21th September 2021.