

To study the composition of minerals and vitamins of some fresh vegetables with special reference to the Kullu District of Himachal Pradesh India.

Estudiar la composición de minerales y vitaminas de algunas verduras frescas con especial referencia al distrito de Kullu de Himachal Pradesh India.

Imtiyaz Rasool Parrey<sup>1,\*</sup>, Tanveera Akhter<sup>1</sup>, Hilal Ul Zaman<sup>2</sup>

1- Department of Chemistry, Women College anantnag Jammu and Kashmir, India-192101

2- Dr. A.P.J Abdul Kalam University Indore Madhya Pradesh, India-452016.

\* Author for correspondence, email: [imtiyazchemistry@gmail.com](mailto:imtiyazchemistry@gmail.com)

#### ABSTRACT

**Aims:** This comparative study was carried out to evaluate the vitamin-C and mineral content of twenty-two different sorts of nutritious and low-cost fresh vegetables. **Methodology:** Vitamin C, calcium (Ca), magnesium (Mg), sodium (Na), potassium (K), phosphorus (P) contents were determined with the flame emission spectrophotometer. **Results:** Vitamin-C was higher in *Momordica charantia* (87.0 mg/100g) and lower in *Vigna unguiculata* (0.10 mg/100g). Calcium and magnesium of the green leafy vegetables were in the range of 191.0 - 12.0 mg/100g and 210.1- 9.0 mg/100g respectively. Sodium content largely varied from 58.1- 2.1 mg/100g in which *Amaranthus cruentus* content higher and *Cucumis sativus* had the lower content. The highest potassium content was recorded in *Cucurbita pepo* (384.0 mg/100g) and the lowest potassium content was found in *Cucumis sativus* (144.20 mg / 100g). Phosphorus content was ranged between 108.0 -13.0 mg /100 g in all the green leafy vegetables. There were significant correlations between mineral contents. **Conclusion:** These results revealed that selected vegetables contain an appreciable amount of vitamin C and minerals and should be included in diets as a supplement of daily allowance needed by the body.

**Keywords:** Fresh vegetable, Vitamin-C, Mineral composition, Kullu.

## RESUMEN

**Objetivos:** Este estudio comparativo se llevó a cabo para evaluar el contenido de vitamina C y minerales de veintidós tipos diferentes de vegetales frescos nutritivos y de bajo costo. **Metodología:** Se determinó el contenido de vitamina C, calcio (Ca), magnesio (Mg), sodio (Na), potasio (K), fósforo (P) con el espectrofotómetro de emisión de llama. **Resultados:** La vitamina C fue mayor en *Momordica charantia* (87,0 mg/100 g) y menor en *Vigna unguiculata* (0,10 mg/100 g). El calcio y el magnesio de los vegetales de hojas verdes estuvieron en el rango de 191,0 - 12,0 mg/100g y 210,1- 9,0 mg/100g respectivamente. El contenido de sodio varió en gran medida de 58,1 a 2,1 mg/100 g, siendo *Amaranthus cruentus* el contenido más alto y *Cucumis sativus* el contenido más bajo. El mayor contenido de potasio se registró en *Cucurbita pepo* (384,0 mg/100g) y el menor contenido de potasio se encontró en *Cucumis sativus* (144,20 mg/100g). El contenido de fósforo osciló entre 108,0 -13,0 mg/100 g en todas las hortalizas de hoja verde. Hubo correlaciones significativas entre los contenidos de minerales. **Conclusión:** Estos resultados revelaron que las verduras seleccionadas contienen una cantidad apreciable de vitamina C y minerales y deben ser incluidas en las dietas como complemento de la cantidad diaria que necesita el organismo.

**Palabra claves:** Verdura fresca, Vitamina C, Composición mineral, Kullu.

## INTRODUCTION

Vegetables play an important role in human nutrition. They are made up of cellulose, hemicellulose and pectin substances that give them their texture and firmness. They provide an adequate amount of dietary fibers, minerals, vitamins and other nutrients, which are usually in short supply in daily diets. They are very important protective foods and useful for the maintenance of health and prevention of various diseases Mohammed et al. (2011). Leafy vegetables have low energy densities thus recommended for weight management. The availability of indigenous vegetables has declined drastically because of the excessive cultivation of field crops. There is also growing ignorance among young people about the existence of these readily available nutritionally rich food plants Nwanekezie EC, et.al. (2014). the supply of minerals is inadequate to meet the dietary requirements of rapidly growing human population in the world. Humans and animals cannot synthesize minerals thus, they must be provided through food and water Mohammed et. al.(2011). Vegetables contain numerous minerals such as Ca, Fe, Cu, P, Zn, Cl, and Na which are vital for growth and

metabolism. The predominant elements found in green leafy vegetables are Ca, K, Fe and Na. These provide an alkalizing effect on acidity produced by other foods, especially those of animal origin Angela C, Rodica C, et.al. 2010. Vitamins are important for human health and among the vitamins, vitamin-C is an essential micronutrient required for normal metabolic functions of the body. Vitamin-C plays an important role as a component of enzymes involved in the synthesis of collagen and carnitine. It is the major water-soluble antioxidant in the human body. Not only does a vitamin-C intake markedly reduce the severity of a cold, but it also effectively prevents secondary viral or bacterial complications. Numerous studies have shown that an adequate intake of vitamin-C is effective in lowering the risk of developing cancers in breast, cervix, colon, rectum, lung, mouth, prostate and stomach. for maintaining good health and to prevent colds, some scientists are of the view that the human body should be kept saturated with vitamin-C Rahman KMM, Rahman MM, et al (2006). Therefore, this study was carried out to evaluate the vitamin and mineral constituents of twenty-two different species of vegetables commonly grown and consumed at Kullu district in Himachal paradise India.

## MATERIAL AND METHODS

Description of study area: The fresh vegetable samples were collected from kullu district Himachal Paradise India (Fig. 1).



Fig. 1: Map showing the study area

Sample collection and preparation: Twenty-two different species of healthy, diseases free, commonly consumed vegetables were collected from farmer's field and home gardens (Table 1). Edible parts of the plants were separated and washed thoroughly under running tap water and then were rinsed in the distilled water. Then the vegetables were cut into pieces and sundry, later then dried in an oven at 70-80°C for 48 hours. The dried portions were grounded into powder using mortar and pestle. Each powdered sample was digested as reported by Anjorin et al. (2010). Briefly, 10g fresh sample was digested with 10-15 ml di-acid mixture (HNO<sub>3</sub>: HClO<sub>4</sub> = 2:1) and heated at 180-200°C on an electric hot plate until white fumes were evolved from the conical flask. The solution was filtered with Whatman filter paper no. 42 and volume made up to 100 ml with distilled water. The aliquot was used separately to determine the mineral contents. The digests were analyzed for the mineral using Buck Scientific model 210VGP Atomic Absorption Spectrophotometer.

Table 1: List of vegetable sample used in this study.

SL. No.	English Name	Scientific Name	Parts used for consumption
1	Bitter gourd	<i>Momordica charantia</i>	Fruits
2	Tomato	<i>Solanum lycopersicum</i>	Fruits
3	Brinjal	<i>Solanum melongena</i>	Fruits
4	Lady's finger	<i>Abelmoschus esculentus</i>	Fruits
5	Potato	<i>Solanum tuberosum</i>	Fruits
6	Bottle gourd	<i>Lagenaria siceraria</i>	Fruits
7	Pumpkin	<i>Cucurbita pepo</i>	Fruits
8	Ash gourd	<i>Benincasa hispida</i>	Fruits
9	Cowpea	<i>Vigna unguiculata</i>	Fruits and seeds
10	Cucumber	<i>Cucumis sativus</i>	Fruits
11	Drum stick	<i>Moringa oleifera</i>	Fruits
12	Pointed gourd	<i>Trichosanthes dioica</i>	Whole part
13	Ridge gourd	<i>Luffa acutangula</i>	Fruits
14	Snake gourd	<i>Trichosanthes cucumerina</i>	Fruits
15	Teasel gourd	<i>Momordica dioica</i>	Fruits and seeds
16	Radish	<i>Raphanus sativus</i>	Whole part
17	Plantain	<i>Musa paradisiaca</i>	Fruits
18	Pea	<i>Pisum sativum</i>	Seeds
19	Bean	<i>Phaseolus vulgaris</i>	Fruits
20	Papaya	<i>Carica papaya</i>	Fruits
21	Chilli	<i>Capsicum frutescens</i>	Fruits
22	Red Amaranth	<i>Amaranthus cruentus</i>	Whole part

Determination of Vitamin-C and minerals: The ascorbic acid content was determined according to Ranganna, 1979. Calcium (Ca) and magnesium (Mg) were determined by the complexometric method of titration using Na<sub>2</sub>-EDTA as a complexing agent Page AL, Miller RH et al (2016) Phosphorus (P) was determined using ascorbic acid as a reductant for color development and reading was recorded with the help of spectrophotometer. Sodium (Na) and potassium (K) content were determined by following the method as described by Terrab et al., (2004) with the help of flame emission spectrophotometer (Spectrolab analytical UK) using appropriate filters.

Statistical analysis: Statistical analyses were carried out by using Statistical Package for Social Science (SPSS) for Windows version 20.0. Results were expressed as mean value  $\pm$  standard deviation (SD). Pearson Correlations was adopted to indicate the measure of correlation and strength of the relationship between variables. For dividing the vegetable species into several groups, cluster analysis (CA) with dendrogram using nearest neighbor method was adopted by using the overall vegetable vitamin-C and mineral content. Other calculations were done by Microsoft excel 2013.

## RESULTS AND DISCUSSION

Fresh vegetables are important sources of nourishment and a vital ingredient in healthy and balanced diets. The roles of vegetables in the maintenance of good health are well known and highly recommended in any diet. Vitamin-C, minerals compositions of the collected vegetables like calcium (Ca), magnesium (Mg), sodium (Na), potassium (K) and phosphorous (P) were presented in Table 2.

Content of Vitamin-C: In this study, the concentration of vitamin-C was between  $87 \pm 0.85$  to  $0.10$  mg/100g. The highest vitamin-C content of  $87 \pm 0.85$  mg/100g was found in *Momordica charantia* followed by Rahman et al. (88 mg/100g). Almost similar content of  $79.50 \pm 0.36$  mg/100g in *Capsicum annum*, which was agreed with Olatunji and Afolayan, 80.6 mg/100g. Although, Emmanuel-Ikpeme et al. found vitamin-C content was  $47.55 \pm 0.44\%$ . According to Table 2, *Vigna unguiculata* contains the least amount of vitamin-C content of  $0.10 \pm 0.00$  mg/100g. A little bit higher vitamin-C content was found by Chikwendu et al. ( $0.59 \pm 0.01$  mg/100g) in husks of cowpea where Soris and Mohan et al observed a much higher ( $34.44 \pm 1.28$  mg/100g) value than the present study.

Content of Calcium (Ca): Calcium among the various vegetables was found to be concentrated in *Amaranthus cruentus* ( $191 \pm 0.8$  mg/100g) than all other vegetable samples, followed by *Abelmoschus esculentus* ( $85 \pm 0.53$  mg/100g), *Phaseolus vulgaris* ( $69.30 \pm 0.26$  mg/100g). *Cucumis sativus* was having the least calcium content of  $12 \pm 0.36$  mg/100g compared to *Solanum tuberosum* and *Solanum lycopersicum* with  $14 \pm 0.10$  and  $14 \pm 0.42$  mg/100g respectively (Table 2). Various values have been previously reported for *Amaranthus cruentus*, Valcárcel-Yamani and Lannes (2017) found ranged at 180.1 - 217.0 mg/100g in amaranth seeds, Topwal found Calcium ranged at 160 - 212 mg/100g, Soriano-García et al. (2018) as 159 mg/100g. In case of *Abelmoschus esculentus*, Habtamu et al. (2017) Found mean calcium content was 81.77 mg/100g which was similar to the present study and higher than those reported for okra seed by Ndangui et al. as 78.65 mg/100g, but lower than those reported as 112.50 - 345.83 mg/100g. In this study, the lowest calcium content found in *Cucumis sativus* ( $12 \pm 0.36$  mg/100g) which was less than Abbey et al. as 20.45 to 25.47 mg/100g.

Content of Magnesium (Mg): Magnesium is important for many processes in the body. Soft tissue magnesium functions as a co-factor of many enzymes involved in energy metabolism, protein synthesis, RNA and DNA synthesis and maintenance of the electrical potential of nervous tissues and cell membranes Pasternak K, Kocot J et al (2010). As shown in Table 2, Amaranthus cruentus contained the largest amount of magnesium ( $210.1 \pm .29$  mg/100g) among the studied vegetables, followed by Carica papaya ( $54 \pm .66$  mg/100g), Phaseolus vulgaris ( $49.50 \pm .30$  mg/100g), Pisum sativum ( $45.40 \pm .43$  mg/100g). In Amaranthus Cruentus, authors found magnesium as 248 mg/100g Soriano-García M, EF, et al. (2018). Valcárcel-Yamani and Lannes found ranged at 279.2 - 319.0 mg/100g which was higher than the present study. Okon et al. (2014) found  $22.33 \pm 0.88$  mg/100g in raw Carica papaya, Ali et al.(2018) as ranged between 10 to 33 mg/100g which was contrasting with present research. In this study, Solanum lycopersicum contains the lowest amount of magnesium  $9 \pm .45$  mg/100g. This value was similar to the findings of Anjum et al. (2013) As ranged between 9.73 to 31.5 mg/100g, Suárez et al. as ranged between 7 to 13 mg/100g. But lower than previous data stated by some authors Nour et al. (2015) as 13.77 to 32.58 mg/100g and Costa et al. as 12 to 20 mg/100g.

Table 2: Vitamin C and Minerals composition of fresh vegetables (mg/100g)

English Name		Ca	Mg	Na	K	P	Vit-C
	Range	25 ± .46	33 ± .44	37 ± 0.72	174 ± .46	22 ± .36	87 ± 0.85
Tomato	Mean ± SD	14 ± .42	9 ± .45	8 ± .56	158 ± .36	26 ± 0.26	20.1 ± .20
	Range	14.50-15.50	8.50-9.40	7.50-8.60	157.60-158.30	25.80-26.30	19.90-20.30
Brinjal	Mean ± SD	19 ± .26	23 ± .26	7 ± .79	172 ± .66	48 ± .26	11.50 ± .26
	Range	18.80-19.30	22.70-23.20	6.40-7.90	171.40-172.70	47.80-48.30	11.20-11.70
Lady's finger	Mean ± SD	85 ± .53	27 ± .26	39 ± .56	181 ± .26	29 ± .56	22.5 ± .30
	Range	84.60-85.60	26.70-27.20	38.50-39.60	180.70-181.20	28.50-29.60	22.20-22.80
Potato	Mean ± SD	14 ± .10	19 ± .66	18 ± .36	285 ± .66	43 ± .26	10.40 ± .43
	Range	13.90-14.10	18.40-19.70	17.70-18.40	284.40-285.70	42.80-43.30	10.10-10.90
Bottle gourd	Mean ± SD	26.83 ± .75	12 ± .26	38 ± .36	150 ± .52	53 ± .26	10.60 ± .44
	Range	26.40-27.70	11.80-12.30	37.70-38.40	149.60-150.60	52.70-53.20	10.30-11.10
Pumpkin	Mean ± SD	32 ± .56	38 ± .26	11 ± .10	384 ± .56	91 ± .66	7.3 ± .26
	Range	31.50-32.60	37.70-38.20	10.90-11.10	437.50-438.60	90.40-91.70	7.10-7.60
Ash gourd	Mean ± SD	28.10 ± .53	16.10 ± .44	38.1 ± .53	148 ± .15	13 ± .36	30.10 ± .72

	Range	27.70-28.70	15.80-16.60	37.70-38.70	147.40-148.90	12.70-13.40	29.50-30.90
Cowpea	Mean ± SD	52.2±.36	42.10±.53	19.10±.43	302±.55	48.77±1.0	0.10±.00
	Range	51.90-52.60	41.70-42.70	18.80-19.60	301.50-302.60	47.70-49.70	0.1-0.1
Cucumber	Mean ± SD	12±.36	10.80±.53	2.1±.26	144.20±.53	16.20±.60	6.9±.26
	Range	11.70-12.40	10.40-11.40	1.90-2.40	143.70-144.70	15.80-16.90	6.70-7.20
Drum stick	Mean ± SD	23.1±.53	27.40±.36	41±.36	256±.53	108±.36	67.80±.53
	Range	22.70-23.70	27.10-27.80	40.70-41.40	255.60-256.60	107.70-108.40	67.40-68.40
Pointed gourd	Mean ± SD	15±.55	14.40±.36	27.10±.53	152.85±.26	17.80±.10	18.50±.36
	Range	14.50-15.60	14.10-14.80	26.70-27.70	152.10-153.60	17.70-17.90	18.20-18.90
Ridge gourd	Mean ± SD	19.40±.36	13.80±.10	37.5±.26	148.95±.36	31.20±.36	4.36±.37
	Range	19.10-19.80	13.70-13.90	37.30-37.80	148.10-149.80	30.80-31.50	4.10-4.80
Snake gourd	Mean ± SD	29.10±.53	16.20±.36	38.10±.43	152.10±.53	27±.66	17.80±.70
	Range	28.70-29.70	15.80-16.50	37.80-38.60	151.70-152.70	26.40-27.70	17.30-18.60
Teasel gourd	Mean ± SD	25.40±.36	18.10±.53	51.10±.53	184.10±.53	25.10±.53	70.30±.26
	Range	25.10-25.80	17.70-18.70	50.70-51.70	183.70-184.70	24.70-25.70	70.10-70.60
Radish	Mean ± SD	23.60±.53	13.90±.20	38.60±.53	150.15±.36	21.50±.30	16.70±.20
	Range	23.20-24.20	13.70-14.10	38.20-39.20	149.80-150.50	21.20-21.80	16.50-16.90
Plantain	Mean ± SD	21.30±.26	27.60±.53	3.7±.20	239.20±.36	20.30±.26	7.2±.36
	Range	21.10-21.60	27.20-28.20	3.50-3.90	238.80-239.50	20.10-20.60	6.80-7.50
Pea	Mean ± SD	42±.55	45.40±.43	3.5±.30	242±.66	105.60±.53	7.10±.44
	Range	41.50-42.60	45.10-45.90	3.20-3.80	241.40-242.70	105.20-106.20	6.80-7.60
Bean	Mean ± SD	69.30±.26	49.50±.30	8.6±.53	168±.36	48±.55	8.7±.20
	Range	69.10-69.60	49.20-49.80	8.20-9.20	167.60-168.30	47.50-48.60	8.50-8.90
Papaya	Mean ± SD	14.5±.26	54±.66	6.10±.53	148.20±.44	14±.55	17.50±.26
	Range	14.30-14.80	53.40-54.70	5.70-6.70	147.80-148.60	13.50-14.60	17.30-17.80
Chilli	Mean ± SD	20.7±.20	42±.36	11±.66	281±.26	29.10±.20	79.50±.36
	Range	20.50-20.90	41.70-42.40	10.40-11.70	280.80-281.30	28.90-29.30	79.20-79.90
Red Amaranth	Mean ± SD	191±0.8	210.1±.29	58.1±.53	260±.53	31.1±.20	41.50±.26



**Content of Sodium (Na):** Sodium is an important mineral and electrolyte necessary for many functions in the body. It has an important role in maintaining water balance within cells and is involved in the proper functioning of both nerve impulses and muscles within the body. Sodium among the various vegetable was found to be concentrated in *Amaranthus cruentus* ( $58.1 \pm 0.53$  mg/100g) than all other vegetable samples, followed by *Momordica dioica* ( $51.10 \pm 0.53$  mg/100g), *Moringa oleifera* ( $41 \pm 0.36$  mg/100g). *Cucumis sativus* was having the least sodium content of  $2.1 \pm 0.26$  mg/100g, compared to *Musa paradisiaca* with  $3.7 \pm 0.20$  mg/100g (Table 2). Different authors reported the level of sodium in *Amaranthus cruentus* was 4 mg/100g which was much lower than the present study Soriano-García M, EF, et al. (2018) Salvi and Katewa (2017) reported sodium as 58 mg/100g in *Momordica dioica* which was similar to this study but in contrast, Kumar et al. found lower content of sodium as  $2.8 \pm 0.2$  mg/100g. The author found the least content of sodium in *Cucumis sativus*, which was similar to the study of Abbey et al. as 2.43 mg/100g, but higher than Olayinka and Etejere (2017) as 0.112 mg/100g. The present found a higher level of Na comparatively than the other studies. It is maybe due to the water salinity of the study area.

**Content of Potassium (K):** Potassium is the most abundant cation in intracellular fluid, where it plays a key role in maintaining cell function, particularly in excitable cells such as muscles and nerves Soriano-García M, EF, et al. (2018). In the present study, Potassium content among the various vegetables was found to be higher in *Cucurbita pepo* ( $384 \pm 0.56$  mg/100g) flesh than all other vegetable samples, followed by *Vigna unguiculata* ( $302 \pm 0.55$  mg/100g), *Capsicum frutescens* ( $281 \pm 0.26$  mg/100g). *Cucumis sativus* was having the least potassium content of  $144.20 \pm 0.53$  mg/100g, compared to *Carica papaya* with  $148.20 \pm 0.44$  mg/100g (Table 2). Various researchers found a variety range of potassium content, Faiku et al. found 172.31 mg/100g in flesh, Elinge et al.  $237.24 \pm 0.09$  mg/100g in the seed of *Cucurbita pepo* which was lower than this study. In contrast, Syed et al. found much higher (809.0 mg/100g) than the present study. Emmanuel-Ikpeme et al. (2015) found  $331.32 \pm 0.02$  mg/100g and Raimi et al. found lower content of potassium ( $98.80 \pm 0.15$  mg/100g) in *Capsicum frutescens*. Abbey et al. (2017) found  $145.55 \pm 0.3$  mg/100g and Labban et al. found  $147.03 \pm 0.25$  mg/100g in *Cucumis sativus* which was similar to this study.

Content of Phosphorus (P): Phosphorus is an essential mineral required for cell structure, signaling, energy transfer, and other important functions. The main function of phosphorus is in the formation of bones and teeth. Phosphorus content among the various vegetables was found to be higher in *Moringa oleifera* ( $108 \pm 0.36$  mg/100g) than all other vegetable samples, followed by *Pisum sativum* ( $105.60 \pm 0.53$  mg/100g), *Cucurbita pepo* ( $91 \pm 0.66$  mg/100g). *Benincasa hispida* was having the least phosphorus content of  $13 \pm 0.36$  mg/100g, compared to *Carica papaya* with  $14 \pm 0.55$  mg/100g (Table 2). Gopalal krishnan et al. (2017) observed the level of phosphorus in *Moringa oleifera* was ranged between 90 to 112 mg/100g that supports the present study. In contrast, Sahay et al. and Abbas et al. (2014) found 70 mg/100g and 70.8 mg/100g respectively. The different authors found a different value of phosphorus in *Cucurbita pepo*. Adebayo et al. found  $118.30 \pm 0.10$  mg/100g which was higher than this study. Elinge et al. (2012) and Hashash et al. (2011) observed lower content of phosphorus as  $47.68 \pm 0.04$  mg/100g and 27.47 mg/100g respectively. Saxena et al. found 19 mg/100g phosphorus in *Benincasa hispida* approximate to this study. The differences in the mineral content of the vegetable plant might be due to soil compositions and the rate of uptake of minerals by an individual plant.

Source analysis of minerals and vitamin-C: Pearson's correlation coefficient was performed to see if some mineral and vitamin-C interrelated with each other and the results are presented in Table 3. The calcium showed significant positive correlation with magnesium ( $r = 0.897^{**}$ ), sodium ( $r = 0.425^*$ ) and vitamin-C ( $r = 0.553^{**}$ ). Similarly, magnesium showed a significant positive correlation with sodium ( $r = 0.776^{**}$ ) and potassium ( $r = 0.422^*$ ). Besides that significant positive correlation was found among sodium-vitamin-C ( $r = 0.435^*$ ) and potassium-phosphate ( $r = 0.574^{**}$ ). Other relationships among the constituents of vegetable samples were not significant. Considering the relationship between the combinations showed a positive significant relationship, which indicates the parameters, was interrelated with each other and maybe originated from the same sources of the study area Abbasi AM, Iqbal J (2013). Furthermore, using the overall nutrient content in vegetable samples, cluster analysis (CA) with dendrogram using nearest neighbor method was adopted to divide the vegetable samples into several groups as shown in Figure 2. Different clusters were formed between different selected vegetables, where the vegetables in each group were of similar growing nature. Moreover, based on nutrient concentrations in some vegetable groups showed strong significant correlations by forming primary groups/clusters with each other (Figure 2).

Table 3: Pearson correlation coefficient matrix for vitamin C and minerals in the vegetables sample.

Pearson Correlations						
	Ca	Mg	Na	K	P	Vit C
Ca	1					
Mg	.897**	1				
Na	.425*	.776**	1			
K	.219	.422*	-.168	1		
P	.038	.048	-.133	.574**	1	
Vit C	.553**	.153	.435*	.056	-.057	1

\*. Correlation is significant at the 0.05 level (2-tailed).  
 \*\*. Correlation is significant at the 0.01 level (2-tailed).




Fig. 2: Cluster analysis of vegetables samples collected from Kullu District Himachal.

As conclusions, this study indicates that there are significant differences in the vitamin-C and mineral composition of different species of fresh vegetables. Vegetables are rich in vitamin-C, minerals such as Ca, Mg, Na, K and P. It contains very low calories, fats but rich in soluble and insoluble fiber at very low cost. Vegetables supply a lot of micronutrients and can fulfill our Recommended Dietary Allowance (RDA). People may be able to know the nutritional status of vegetables through this study. Further study should be conducted on the assessment of different minerals and vitamins in vegetables for food and nutritional security in the study area.

#### REFERENCES

- Abbey BW, Nwachoko N and Ikiroma GN. 2017. Nutritional Value of Cucumber Cultivated in Three Selected States of Nigeria. *Biochem Anal Biochem.*; 6: 328.
- Abbasi AM, Iqbal J, Khan MA, Shah MH 2013. Health risk assessment and multivariate apportionment of trace metals in wild leafy vegetables from Lesser Himalayas, Pakistan. *Ecotoxicol. Environ. Saf.*; 92: 237–244.
- Abbas RK, Elsharbasy FS, Fadlemula AA. 2018. Nutritional Values of *Moringa oleifera*, Total Protein, Amino Acid, Vitamins, Minerals, Carbohydrates, Total Fat and Crude Fiber, under the Semi-Arid Conditions of Sudan. *J Microb Biochem Technol.*; 10: 56-58.
- Adebayo OR, Farombi AG, Oyekanmi AM. Proximate, Mineral and Anti-Nutrient Evaluation of Pumpkin Pulp (*Cucurbita Pepo*). *IOSR Journal of Applied Chemistry*. 2013; 4(5): 25-28.
- Ali A, Devarajan S, Waly MI, Essa MM, Rahman MS 2012. Nutritional and medicinal values of papaya (*Carica papaya* L.). In *Natural Products and Their Active Compounds on Disease Prevention.*; 307-324.
- Angela C, Rodica C, Andrea MZ, Tonea E, Giuchici C. 2010. Chemical Composition of Common Leafy Vegetables. *Studia Universitatis Series of Life Sciences.*; 20(2): 45-48.
- Anjum S, Hamid A, Ghafoor A, Tahira R, Shah SZA, Awan SI, et al. Evaluation of Biochemical Potential in Tomato (*Solanum lycopersicum*) Germplasms. *Pak. J. Agri. Sci.* 2020; 57(1): 177-187.

- Anjorin TS, Ikokoh P, Okolona S. 2010. Mineral Composition of *Moringa Oleifera* leaves, Pods, and seed from two region in Abuja, Nigeria. *International Journal of Agriculture and Boilogy.*; 12: 431 – 434.
- Asaolu SS, Asaolu MF 2010. Trace Metal Distribution in Nigerian Leafy Vegetables. *Pakistan Journal of Nutrition.*; 9(1): 91-92. doi: 10.3923/pjn. 91.92
- Chikwendu JN, Igbatim AC, Obizoba IC. 2014. Chemical Composition of Processed Cowpea Tender Leaves and Husks. *International Journal of Scientific and Research Publications.* 4(5): 1-5.
- Emmanuel-Ikpeme C, Henry P, Okiri OA. 2014. Comparative Evaluation of the Nutritional, Phytochemical and Microbiological Quality of Three Pepper Varieties. *Journal of Food and Nutrition Sciences.*; 2(3): 74-80.
- Elinge CM, Muhammad A, Atiku FA, Itodo AU, Peni IJ, Sanni OM, et al. 2012. Proximate, Mineral and Anti-nutrient Composition of Pumpkin (*Cucurbitapepo L*) Seeds Extract. *International Journal of Plant Research.*; 2(5): 146-150. doi: 10.5923/j.plant.20120205.02.
- Faiku F, Haziri A, Gashi F, Troni N. 2016. Nutrient Composition of *Cucurbita Melo* Growing in Kosovo. *Chemistry. Bulgarian Journal of Science Education.*; 25(3): 438-447.
- Gopalakrishnan L, Doriya K, Kumar DS. *Moringa oleifera*: 2016. A review on nutritive importance and its medicinal application. *Food Science and Human Wellness.*; 5: 49–56.
- Habtamu FG, Haki GD et al. 2018. Nutritional and Antinutritional Evaluation of Indigenous Ethiopian Okra (*Abelmoschus Esculentus*) Seed Accessions. *Afr. J. Food Agric. Nutr. Dev.*; 18(1): 13019-13033.
- Hashash MM, El-Sayed MM, Abdel-Hady AA, Hady HA, Morsi EA. Nutritional Potential, Mineral Composition and Antioxidant Activity Squash (*Cucurbita Pepo L.*) Fruits Grown in Egypt. *European Journal of Biomedical and Pharmaceutical Sciences.* 2017; 4(3): 05-12.
- Kumar GMP, Chikkapaih L and Nagayya S. 2016. Nutritional analysis of edible wild plants used by Hakki Pikki tribes of Hassan district, Karnataka, India. *Int J Pharm Pharm Sci.*; 8(8): 390-393.

- Labban ML, Mouzik JEM, Shekh Otham ADA. 2017. Comparison of Sodium and Potassium Content in Fresh Produce, and their Contribution to the Daily Intake. *J Adv Res Food Sci Nutr.*; 3(3&4): 1-8.
- Mohammed MI, Sharif N. 2011. Mineral Composition of Some Leafy Vegetables consumed in Kano, Nigeria. *Nigerian Journal of Basic and Applied Science.* 19(2): 208-212.
- Ndangui CB, Kimbonguila et al. Nutritive 2010. Composition and Properties Physico-chemical of gumbo (*Abelmoschus esculentus* L.) Seed and Oil. *Research Journal of Envir Earth Sci.*; 2(1): 49-54.
- Nour V, Trandafir I, Ionica ME. 2013. Antioxidant Compounds, Mineral Content and Antioxidant Activity of Several Tomato Cultivars Grown in Southwestern Romania. *Not Bot Horti Agrobo.*; 41(1): 136-142.
- Nwanekezie EC, Obiakor-Okeke PN 2014. Mineral Content of Five Tropical Leafy Vegetables and Effect of Holding Methods and Time. *American Journal of Experimental Agriculture.*; 4(12): 1709-1717.
- Odhav B, Beekrum S, Baijnath H. 2006. Preliminary assessment of nutritional value of traditional leafy vegetables in KwaZulu-Natal, South Africa. *Journal of Food Composition and Analysis.*; 20: 430-435. doi: 10.1016/j.jfca.2006.04.015
- Olatunji TL, Afolayan AJ. 2008. The suitability of chili pepper (*Capsicum annum* L.) for alleviating human micronutrient dietary deficiencies: A review. *Food Sci Nutr.*; 6: 2239-2251.
- Olayinka BU, Etejere EO. 2018. Proximate and Chemical Compositions of Watermelon (*Citrullus lanatus* (Thunb.) Matsum and Nakai cv Red and Cucumber (*Cucumis sativus* L. cv Pipino). *International Food Research Journal.*; 25(3): 1060-1066.
- Okon WI, Ogri AI, Igile GO, Atangwho I 2017. Nutritional quality of raw and processed unripe *Carica papaya* fruit pulp and its contribution to dietary diversity and food security in some peasant communities in Nigeria. *Int. J. Biol. Chem. Sci.*; 11(3): 1000-1011.
- Page AL, Miller RH and Keeney DR. *Methods of soil analysis; 2. Chemical and microbiological properties*, 2. Aufl. 1184 S., American SOC. of Agronomy (Publ.), Madison, Wisconsin, USA.

- Pasternak K, Kocot J and Horecka A 2010. Biochemistry of Magnesium. J. Elementol. 15(3): 601–616.
- Rahman KMM, Rahman MM, Islam MS, Begum SA. 2006. A simple UV-spectrophotometric method for the determination of vitamin C content in various fruits and vegetables at Sylhet area in kullu. Journal of Biological Sciences.; 6(2): 388 – 392.
- Rahman AHMM, Anisuzzaman M, Ahmed F, Islam AKMR, Naderuzzaman ATM. 2008. Study of Nutritive Value and Medicinal Uses of Cultivated Cucurbits. Journal of Applied Sciences Research; 4(5): 555-558.
- Raimi MM, Shittu SA, Oyetade OA. 2014. Physicochemical Properties and Mineral Composition of Capsicum annum and Capsicum frutescens Oils. IOSR Journal of Applied Chemistry.; 7(1): 112-116.
- Ranganna S. Titratable acidity. 1979. Manual of analysis of food and vegetable Products, Tata McGraw-Hill, Delhi, India.
- Sahay S, Yadav U, Srinivasamurthy S 2017. Potential of *Moringa oleifera* as a functional food ingredient- A review. International Journal of Food Science and Nutrition.; 2(5): 31-37.
- Salvi J, Katewa SS. 2015. Nutritional Composition of *Momordica dioica* fruits: As a wild vegetable. J. Food Pharm.Sci.; 3: 18-22.
- Sami R, Alshehry G, Ma Y, Abdelazez A, Benajiba N. 2019. Evaluation of Some Specific Components Existences in Okra (*Abelmoschus Esculentus L. (Moench)*) Cultivated from Different Areas. Journal of Food and Nutrition Research; 7(2): 155-161.
- Saxena PK, Sharma RD, Sharma KK, Ritu G, Sachin T. Another Look On: *Benincasa hispida*. International Journal of Pharmaceutical and Chemical Sciences. 2016; 5(2): 150-156.
- Shammi M, Rahman MM, Bondad SE, Bodrud-Doza M. 2019. Impacts of Salinity Intrusion in Community Health: A Review of Experiences on Drinking Water Sodium from Coastal Areas of kullu Healthcare.; 7: 50.
- Soriano-García M, EF, et al. 2018. Nutritional functional value and therapeutic utilization of Amaranth. J Anal Pharm Res.; 7(5): 596–600.

Soris PT, Mohan VR. 2011. Chemical Analysis and Nutritional Assessment of Two Less Known Pulses of Genus *Vigna*. Tropical and Subtropical Agroecosystems.; 14: 473 – 484.

Suarez MH, Rodriguez EMR, Romero CD. 2007. Mineral and trace element concentrations in cultivars of tomatoes. Food Chemistry.; 104: 489–499.

Syed QA, Akram M, Shukat R. 2019. Nutritional and Therapeutic Importance of the Pumpkin Seeds. Biomed J Sci & Tech Res.; 21(2): 15798-15803.

Terrab A, Hernanz D, Heredia FJ. 1982. Sample treatment procedures for the determination of mineral constituents in honey by Inductively Coupled Plasma Optical Emission Spectrometry. J. Agric. Food Chem. 2004; 52: 3441-3445.

Topwal M. A Review on Amaranth 2019.: Nutraceutical and Virtual Plant for Providing Food Security and Nutrients. Acta Scientific Agriculture.; 3(1): 09-15.

Valcárcel-Yamani B, Lannes SCS. 2012. Applications of Quinoa (*Chenopodium Quinoa* Willd.) and Amaranth (*Amaranthus* Spp.) and Their Influence in the Nutritional Value of Cereal Based Foods. Food and Public Health.; 2(6): 265-275.

Weaver CM. Potassium and health. Adv. Nutr. 2013; 4: 368S–377S.

Witt KA. The Nutrient Content of *Moringa oleifera* Leaves. ECHO Research Note. 2013; 1:1.

Wolosik K, Markowska A. *Amaranthus Cruentus* 2019. Taxonomy, Botanical Description, and Review of its Seed Chemical Composition. Natural Product Communications.; 1–10.

Received: 26<sup>th</sup> Jule 2021; Accepted:30<sup>th</sup> August 2021; First distribution: 03<sup>th</sup> February 2022.